

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07K 14/47, A61K 38/04, G01N 33/68	A1	(11) International Publication Number: WO 96/13517 (43) International Publication Date: 9 May 1996 (09.05.96)
(21) International Application Number: PCT/EP95/04201 (22) International Filing Date: 25 October 1995 (25.10.95) (30) Priority Data: 94203128.7 27 October 1994 (27.10.94) EP (34) Countries for which the regional or international application was filed: AT et al. 95200886.0 7 April 1995 (07.04.95) EP (34) Countries for which the regional or international application was filed: AT et al. (71) Applicant (for all designated States except US): AKZO NOBEL N.V. [NL/NL]; Velperweg 76, NL-6824 BM Arnhem (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): BOOTS, Anna, Maria, Helena [NL/NL]; Verlengde Torenstraat 10, NL-5366 AV Megen (NL). VERHEIJDEN, Gijsbertus, Franciscus, Maria [NL/NL]; Heischouw 7, NL-5345 XT Oss (NL). (74) Agent: HERMANS, F., G., M.; Postbus 20, NL-5340 BH Oss (NL).		(81) Designated States: AU, BR, CA, CN, FI, HU, JP, KR, MX, NO, NZ, PL, RU, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: NOVEL PEPTIDES DERIVED FROM AUTOANTIGEN FOR USE IN IMMUNOTHERAPY OF AUTOIMMUNE DISEASES		
(57) Abstract <p>The present invention relates to novel peptides derived from the autoantigen HC gp-39, said peptides comprising at least one of the amino acid sequences FGRSFTLAS (SEQ ID No.1), FTLASSETG (SEQ ID No. 2), YDDQESVKS (SEQ ID No.3) and FSKIASNTQ (SEQ ID No.4). The peptides resemble MHC Class II restricted T-cell epitopes present on the autoantigen HC gp-39 in articular cartilage. HC gp-39 and said peptides can be used in antigen-specific treatment of articular cartilage destruction in autoimmune diseases to induce tolerance of the immune system. The autoantigen HC gp-39 and said peptides are also suitable to induce arthritis in non-human animals, preferably mice. The invention furthermore relates to pharmaceutical compositions comprising said autoantigen and/or said peptides, a diagnostic method for the detection of autoreactive T cells in a test sample and test kits to be used in said method.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

NOVEL PEPTIDES DERIVED FROM AUTOANTIGEN FOR USE IN IMMUNOTHERAPY OF AUTOIMMUNE DISEASES

5 The invention relates to a novel autoantigen and peptides derived thereof, their use in treatment of chronic destruction of articular cartilage in autoimmune diseases, pharmaceutical compositions comprising said peptides, a diagnostic method for the detection of autoreactive T cells in a test sample and test kits to be used in said method.

10 The immune system is established on a principle of discrimination between foreign antigens (non-self antigens) and autoantigens (self antigens, derived from the individuals own body) achieved by a build in tolerance against the autoantigens

15 The immune system protects individuals against foreign antigens and responds to exposure to a foreign antigen by activating specific cells such as T- and B lymphocytes and producing soluble factors like interleukins, antibodies and complement factors. The antigen to which the immune system responds is degraded by the antigen presenting cells (APCs) and a fragment of the antigen is expressed on the cell surface associated with a major histocompatibility complex (MHC) class II glycoprotein. The MHC-glycoprotein-antigen-fragment complex is presented to a T cell which by virtue of its T cell receptor recognizes the
20 antigen fragment conjointly with the MHC class II protein to which it is bound. The T cell becomes activated, i.e. proliferates and/or produces interleukins, resulting in the expansion of the activated lymphocytes directed to the antigen under attack (Grey et al., Sci. Am., 261 38-46, 1989).

25 Self antigens are also continuously processed and presented as antigen fragments by the MHC glycoproteins to T cells (Jardetsky et al., Nature 353 326-329, 1991). Self recognition thus is intrinsic to the immune system. Under normal circumstances the immune system is tolerant to self antigens and activation of the immune response by these self antigens is avoided.

30 When tolerance to self antigens is lost, the immune system becomes activated against one or more self antigens, resulting in the activation of autoreactive T cells and the production of autoantibodies. This phenomenon is referred to as autoimmunity. As the immune response in

general is destructive, i.e. meant to destroy the invasive foreign antigen, autoimmune responses can cause destruction of the body's own tissue

5 The contribution of T cells to autoimmune diseases has been established by several studies. In mice, experimental autoimmune encephalomyelitis (EAE) is mediated by a highly restricted group of T cells, linked by their specificity for a single epitope of myelin basic protein (MBP) complexed to an MHC class II molecule. In the Lewis rat, a species with high susceptibility to various autoimmune diseases, disease has been shown to be mediated by T cells. In humans autoimmune diseases are also thought to be associated with the development of auto-
10 aggressive T cells

A destructive autoimmune response has been implicated in various diseases such as rheumatoid arthritis (RA), in which the integrity of articular cartilage is destroyed by a chronic inflammatory process resulting from the presence of large numbers of activated lymphocytes and
15 MHC class II expressing cells. The mere presence of cartilage appears necessary for sustaining the local inflammatory response. It has been shown that cartilage degradation is associated with the activity of cartilage-responsive autoreactive T cells in RA (Sigall et al., Clin. Exp. Rheumatol. 6:59, 1988; Glant et al., Biochem. Soc. Trans. 18:796, 1990; Burmester et al., Rheumatoid arthritis. Smolen, Kalden, Maini (Eds) Springer-Verlag Berlin Heidelberg, 1992). Furthermore, removal of cartilage from RA patients by surgery was shown to reduce the inflammatory process
20 (G.S. Panayi et al., Clin. Exp. Rheumatol. 11(suppl 8): S1-S8, 1993). The cartilage proteins are therefore considered to be target autoantigens which are competent of stimulating T cells. Activation of these autoreactive T cells leads to development of autoimmune disease.

25 The inflammatory response resulting in the destruction of the cartilage can be treated by several drugs, such as for example steroid drugs. However, these drugs are often immunosuppressive drugs that are nonspecific and have toxic side effects. The disadvantages of nonspecific immunosuppression makes this a highly unfavourable therapy.

30 The antigen-specific, nontoxic immunosuppression therapy provides a very attractive alternative for the nonspecific immunosuppression. This antigen-specific therapy involves the treatment of patients with the target autoantigen or with synthetic T cell-reactive peptides.

- 3 -

derived from the autoantigen. These synthetic peptides correspond to T cell epitopes of the autoantigen and can be used to induce specific T cell tolerance both to themselves and to the autoantigen. Although it seems paradoxical to desensitize the immune system with the very same antigen responsible for activating the immune system, the controlled administration of the target (auto)antigen can be very effective in desensitization of the immune system

To effectively use the tolerance therapy to treat the T cell mediated cartilage destruction, there is a great need to identify the responsible autoantigen and to find T cell-reactive peptides which can desensitize patients against the autoantigen that is activating the T cells responsible for the inflammatory process.

It is an object of the invention to provide the autoantigen and T cell reactive peptides derived from said autoantigen which are able to induce specific T cell tolerance to the responsible cartilage antigen in patients suffering from T cell-mediated cartilage destruction. It is another object of the invention to provide a method for detecting autoreactive T cells involved in the destruction of articular cartilage and test kits to be used in said method

It was surprisingly found that Human Cartilage glycoprotein 39 (herein after referred to as HC gp-39) is a target autoantigen in RA patients which activates specific T cells, thus causing or mediating the inflammatory process. HC gp-39 derived peptides were predominantly recognized by autoreactive T cells from RA patients but rarely by T cells from healthy donors, thus indicating that HC gp-39 is an autoantigen in RA. The arthritogenic nature of HC gp-39 was further substantiated in the Balb/c mouse. A single, subcutaneous injection of said protein in Balb/c mice was able to initiate arthritic signs in the animals. The course of the HC gp-39-induced disease was characterized by relapses occurring periodically in fore paws and/or hind paws and gradually developed from a mild arthritis into a more severe form. Also, a symmetrical distribution of afflicted joints was observed which is, together with the observation of recurrent relapses and nodule formation, reminiscent of disease progression in arthritis, especially RA.

Even more surprisingly it was found that administration of HC gp-39 resulted in immunological tolerance and, more importantly, in delayed and/or suppressed arthritic development

5 HC gp-39 is present in serum of both patients and healthy adults, although the serum concentration of the protein is about twice as much in patients as compared to healthy adults. Furthermore, mRNA coding for HC gp-39 can be found in synovial specimens or cartilage obtained from RA patients, whereas cartilage of healthy adults, obtained at surgery, does not contain a significant amount of said mRNA. When articular chondrocytes and synovial cells are cultured, their major secretory product becomes HC gp-39 (Hakala et al., J Biol Chem., Vol 268, 34 25803, 1993). The arthritogenic nature of HC gp-39 was neither described nor suggested in the Hakala et al publication, nor in any other publication.

10 A further object of the invention is achieved by peptides comprising a subsequence of the autoantigen HC gp-39, characterized in that said peptides comprise one or more of the amino acid sequences FGRSFTLAS (SEQ ID No 1), FTLASSETG (SEQ ID No 2), YDDQESVK (SEQ ID No 3) and FSKIASNTQ (SEQ ID No 4).

15 More specifically, a peptide according to the invention comprises one or more of the amino acid sequences PTFGRSFTLASSE (SEQ ID No 5), RSFTLASSETGVG (SEQ ID No 6), VGYDDQESVSKV (SEQ ID No 7) and SQRFSKIASNTQSR (SEQ ID No 8).

20 "Subsequence" is understood to be defined as "a part" and should not be mistaken to encompass the entire protein. Preferably the peptides according to the invention have an amino acid sequence of 9-55 amino acid residues. More preferably the peptides according to the invention have an amino acid sequence of 9-35, in particular 9-25 amino acid residues. Much more preferred are peptides having an amino acid sequence of 9-15 amino acid residues. Highly preferred are peptides having an amino acid sequence of 13 or 14 amino acid residues, such as for example peptides having the amino acid sequences given in SEQ ID No 5-8.

25 Multimers of the peptide according to the invention, such as for example a dimer or a trimer, in which the monomer sequences optionally can be separated by spacer residues are also within the scope of the invention. Such multimers provide a multitude of the T cell epitopes given in SEQ ID No 's 1-8.

30 In the peptides according to the invention the amino acid sequences given in SEQ ID No 1-8 can be flanked by flanking regions which may correspond to the native flanking sites of the corresponding amino acids in the amino acid sequence of the HC gp-39 protein or other

proteins in which said amino acid sequences are present. Alternatively, said flanking regions can also be non-native amino acid sequences made up of random amino acid residues. These non-native flanking sites can be used to stabilize the peptides, thus increasing their biological availability. To increase the biological availability of the peptides according to the invention,
5 non-native flanking sites are preferred.

The amino acid sequences given in SEQ ID No.'s 1-4, more specifically the sequences given in SEQ ID No.'s 5-8 resemble MHC class II restricted T cell epitopes which are present on HC gp-39. MHC class II restricted T-cell epitopes on HC gp-39 are displayed by the regions
10 103-116, 259-271, 263-275 and 326-338 of the amino acid sequence of HC gp-39 (starting from the methionine in the signal sequence, see Hakala et al. 1993). Thus, according to the invention, the peptides can also be understood to encompass fragments of the autoantigen HC gp-39 which comprise one or more of the above identified MHC Class II restricted T-cell epitopes and they are also within the scope of the invention.

15 The peptides according to the invention are T-cell reactive peptides, which are recognized by and are able to stimulate activated, autoreactive T-cells. These autoreactive T-cells are found in the blood of RA patients but rarely in healthy donors.

Thus, according to the invention HC gp-39 protein or the synthetic peptides, said
20 peptides resembling the MHC Class II restricted T-cell epitopes present on the target autoantigen HC gp-39, are very suitable for use in a therapy to induce specific T-cell tolerance to HC gp-39 in patients suffering from T-cell mediated cartilage destruction, such as for example arthritis, more specifically rheumatoid arthritis.

WO 95/01995 and WO 95/02188 describe the diagnostic use of HC gp-39 as a marker
25 for RA. The arthritogenic nature of HC gp-39 is neither disclosed nor suggested. Nowhere do they hint or suggest towards the use of HC gp-39, fragments thereof or T-cell reactive peptides according to the present invention in the antigen or peptide specific therapy to induce T-cell specific tolerance to the HC gp-39 in the cartilage under attack.

The preparation of the peptides according to the invention is effected by means of one of the known organic chemical methods for peptide synthesis HC gp-39 and the peptides can also be prepared with the aid of recombinant DNA techniques

5 The organic chemical methods for peptide synthesis are considered to include the coupling of the required amino acids by means of a condensation reaction, either in homogeneous phase or with the aid of a so-called solid phase.

The condensation reaction can be carried out as follows:

10 a) condensation of a compound (amino acid, peptide) with a free carboxyl group and protected other reactive groups with a compound (amino acid, peptide) with a free amino group and protected other reactive groups, in the presence of a condensation agent.

 b) condensation of a compound (amino acid, peptide) with an activated carboxyl group and free or protected other reaction groups with a compound (amino acid, peptide) with a free amino group and free or protected other reactive groups

15 Activation of the carboxyl group can take place, inter alia, by converting the carboxyl group to an acid halide, azide, anhydride, imidazolide or an activated ester, such as the N-hydroxy-succinimide, N-hydroxy-benzotriazole, p-nitrophenyl ester, N-hydroxy-benzotriazole ester or pentafluorophenol ester.

20 The most common methods for the above condensation reactions are the carbodiimide method, the azide method, the mixed anhydride method and the method using activated esters, such as described in The Peptides, Analysis, Synthesis, Biology Vol 1-3 (Ed. Gross, E. and Meienhofer, J.) 1979, 1980, 1981 (Academic Press, Inc.)

25 Preparation of suitable fragments of above-mentioned peptides according to the invention using the "solid phase" is for instance described in J. Amer. Chem. Soc. 85 2149 (1963) and Int. J. Peptide Protein Res. 35 161-214 (1990). The coupling of the amino acids of the peptide to be prepared usually starts from the carboxyl end side. For this method a solid phase is needed on which there are reactive groups or on which such groups can be introduced. This can be, for
30 example, a copolymer of benzene and divinylbenzene with reactive chloromethyl groups, or a polymeric solid phase rendered reactive with hydroxymethyl or amine-function.

A particularly suitable solid phase is, for example, the p-alkoxybenzyl alcohol resin (4-hydroxy-methyl-phenoxy-methyl-copolystyrene-1% divinylbenzene resin), described by Wang (1974) J. Am. Chem. Soc. 95 1328. After synthesis the peptides can be split from this solid phase under mild conditions.

5 After synthesis of the desired amino acid sequence, detaching of the peptide from the resin follows, for example, with trifluoroacetic acid, containing scavengers, for example triisopropyl silane, anisole or ethanedithiol, thioanisole.

10 The reactive groups which may not participate in the condensation reaction are, as stated, effectively protected by groups which can be removed again very easily by hydrolysis with the aid of acid, base or reduction. Thus, a carboxyl group can be effectively protected by, for example, esterification with methanol, ethanol, tertiary butanol, benzyl alcohol or p-nitrobenzyl alcohol and amines linked to solid support.

15 Groups which can effectively protect an amino group are the ethoxycarbonyl, benzyloxycarbonyl, t-butoxy-carbonyl (t-boc) or p-methoxy-benzyloxycarbonyl group, or an acid group derived from a sulphonic acid, such as the benzene-sulphonyl or p-toluene-sulphonyl group, but other groups can also be used, such as substituted or unsubstituted aryl or aralkyl groups, for example benzyl and triphenylmethyl, or groups such as ortho-nitrophenyl-sulphenyl
20 and 2-benzoyl-1-methyl-vinyl. A particularly suitable α -amino-protective group is, for example, the base-sensitive 9-fluorenyl-methoxycarbonyl (Fmoc) group [Carpino & Han (1970) J. Amer. Chem. Soc. 92 5748].

25 A more extensive account of possible protecting groups can be found in The Peptides, Analysis, Synthesis, Biology, Vol. 1 - 9 (Eds. Gross, Udénfriend and Meienhofer) 1979 - 1987 (Academic Press, Inc.).

30 The protective groups can be split off by various conventional methods, depending on the nature of the particular group, for example with the aid of trifluoroacetic acid or by mild reduction, for example with hydrogen and a catalyst, such as palladium, or with HBr in glacial acetic acid.

As already indicated above, HC gp-39 and the peptides according to the invention can likewise be prepared with the aid of recombinant DNA techniques. For this purpose, a nucleic acid sequence which codes for HC gp-39 or a peptide according to the invention or a multimer of said peptide is inserted into an expression vector. Suitable expression vectors are, amongst
5 others, plasmids, cosmids, viruses and YAC's (Yeast Artificial Chromosomes) which comprise the necessary control regions for replication and expression. The expression vector can be brought to expression in a host cell. Suitable host cells are, for instance, bacteria, yeast cells and mammalian cells. Such techniques are well known in the art (Sambrook et al., Molecular
10 Cloning a Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, 1989)

Although it seems paradoxical to desensitize the immune system with the very same antigen responsible for activating the immune system, the controlled administration of HC gp-39
15 and/or peptides comprising a subsequence of HC gp-39 can be effective in desensitization of the immune system. According to the invention, patients in which the cartilage is under attack of autoreponsive T cells can be treated with a pharmaceutical composition comprising HC gp-39, or one or more peptides according to the invention and a pharmaceutical acceptable carrier in order to make the specific autoreactive T cells of these patients tolerant to the HC gp-39 in the
20 cartilage under attack and other self antigens carrying the identified T cell epitopes having one of the amino acid sequences given in SEQ ID No. 1-8 and to diminish the inflammatory response. Very suitable peptides to be used in a pharmaceutical composition according to the invention are the peptides having an amino acid sequence given in SEQ ID No. 5, 6, 7 and 8.

Also very suitable to be used in a pharmaceutical composition according to the invention
25 are DNA (expression)vectors comprising DNA which encodes for HC gp-39 or one or more of the peptides according to the invention. Upon delivery the DNA (expression)vector can provide by expression a level of the recombinant HC gp-39 protein or peptides according to the invention which is similar to the level which would be achieved by direct administration of a pharmaceutical composition comprising the HC gp-39 protein or peptides.

30 The autoantigen and peptides according to the invention have the advantage that they have a specific tolerizing effect on the autoreactive T cells thus leaving the other components of

the immune system intact as compared to the nonspecific suppressive effect of the immunosuppressive steroid drugs. Treatment with the autoantigen or peptides according to the invention will be safe and no toxic side effects will occur.

Tolerance can be attained by administering high or low doses of the autoantigen or peptides according to the invention. The amount of autoantigen or peptide will depend on the route of administration, the time of administration, the age of the patient as well as general health conditions and diet.

In general, a dosage of 0.01 to 1000 μg of peptide or protein per kg body weight, preferably 0.5 to 500 μg , more preferably 0.1 to 100 μg of peptide or protein can be used.

Pharmaceutical acceptable carriers are well known to those skilled in the art and include, for example, sterile saline, lactose, sucrose, calcium phosphate, gelatin, dextrin, agar, pectin, peanut oil, olive oil, sesame oil and water. Other carriers may be, for example MHC class II molecules, if desired embedded in liposomes.

In addition the pharmaceutical composition according to the invention may comprise one or more adjuvants. Suitable adjuvants include, amongst others, aluminium hydroxide, aluminium phosphate, amphigen, tocopherols, monophosphoryl lipid A, muramyl dipeptide and saponins such as Quil A. The amount of adjuvant depends on the nature of the adjuvant itself.

Furthermore the pharmaceutical composition according to the invention may comprise one or more stabilizers such as, for example, carbohydrates including sorbitol, mannitol, starch, sucrose-dextrin and glucose, proteins such as albumin or casein, and buffers like alkaline phosphates.

Suitable administration routes are intramuscular injections, subcutaneous injections, intravenous injections or intraperitoneal injections, oral and intranasal administration. Oral and intranasal administration are preferred administration routes.

Due to its arthritogenic nature, HC gp-39 or the peptides according to the invention can be used to induce clinical arthritis in non-human mammals. Upon administration of small amounts of HC gp-39 or one or more of the peptides according to the invention, arthritic signs will develop in said mammals resulting in a disease pattern reminiscent of disease progression in arthritis, especially rheumatoid arthritis. When Balb/c mice were injected subcutaneously with HC gp-39 protein, the animals developed arthritic signs. The course of the HC gp-39-induced

disease was characterized by relapses occurring periodically in fore paws and/or hind paws and gradually developed from a mild arthritis into a more severe form. Also, a symmetrical distribution of afflicted joints was observed which is, together with the observation of recurrent relapses and nodule formation, reminiscent of disease progression in arthritis, especially RA.

5 Thus, these afflicted animals provide an adequate animal model to study the mechanism underlying the initiation and progression of arthritic development. Additionally, said afflicted animals can be used to search for new drugs to treat arthritis and to study the effect of these drugs upon the arthritic development. Preferably mice are used as animal model for arthritis, especially rheumatoid arthritis.

10 To induce arthritis in said mammals, suitable amounts of HC gp-39 or one or more of the peptides according to the invention have to be administered. Suitable amounts are 0.1-1000 µg, preferably 1-100 µg, more preferably 10-50 µg per kg body weight. The amount of HC gp-39 or the peptides will depend on the route of administration, time of administration and the type of animal used. Suitable administration routes are the same as described before. To induce the
15 effect of arthritis induction, the HC gp-39 protein or peptides according to the invention may comprise one or more stabilizers or adjuvants as described before.

HC gp-39 or the peptides according to the invention are also very suitable for use in a diagnostic method to detect the presence of activated autoreactive T cells involved in the chronic
20 inflammation of the articular cartilage.

The diagnostic method according to the invention comprises the following steps:

a) isolation of the peripheral blood mononuclear cells (PBMC) from a blood sample of an individual,

b) culture said PBMC under suitable conditions,

25 c) incubation of said PBMC culture in the presence of the autoantigen or one or more peptides derived thereof according to the invention, and

d) detection of a response of T cells, for example a proliferative response, indicating the presence of activated autoreactive T cells in the individual.

In case of detection of a response by measuring the proliferative response of the
30 autoreactive T cells, the incorporation of a radioisotope such as for example ³H-thymidine is a measure for the proliferation. A response of the autoreactive T cells present in the PBMC can

also be detected by measuring the cytokine release with cytokine-specific ELISA, or the cytotoxicity with $^{51}\text{Chromium}$ release. Another detection method is the measurement of expression of activation markers by FACS analysis, for example of IL-2R. A diagnostic composition comprising one or more of the peptides according to the invention and a suitable
5 detecting agent thus forms part of the invention. Depending on the type of detection, the detection agent can be a radioisotope, an enzyme, or antibodies specific for cell surface or activation markers.

Also within the scope of the invention are test kits which comprise one or more peptides according to the invention. These test kits are suitable for use in a diagnostic method according
10 to the invention.

Thus, the present invention provides for a method to detect whether autoaggressive T cells reactive towards HC gp-39 are present in patients suffering from T-cell mediated cartilage destruction such as for example arthritis, in particular rheumatoid arthritis. If HC gp-39-specific T cells are present, tolerization of these T cells with a pharmaceutical composition comprising
15 HC gp-39 or peptides according to the present invention or combinations thereof can delay or suppress arthritis development.

The following examples are illustrative for the invention and should in no way be interpreted as limiting the scope of the invention.

LEGENDS TO THE FIGURES

Figure 1: Initiation and progression of arthritis in HC gp-39 tolerized and non-tolerized Balb/c mice. The total arthritic score of the afflicted animals per day following sensitization is
25 given for both fore- and hind paw. The number of afflicted animals per day following sensitization is also given.

EXAMPLE 1

METHODS

Patients

5 Peripheral blood mononuclear cells (PBMC) from patients who were diagnosed as suffering from RA according to the American Rheumatism Association (ARA) criteria (Arnett et al., Arthritis Rheum 31 315, 1988) were collected. The severity of disease of RA patients ranged from stage I - IV as determined by Rontgenscore.

10 PBMC's from healthy donors carrying the DR4Dw4 (DRB1*0401) or the DR1 specificity were collected as a control. Also PBMC's from two healthy donors who did not carry one of the RA associated DR molecules were collected.

MHC typing

15 Patient and healthy donor PBMC chromosomal DNA extracts, were analysed using the Dynal DR 'low resolution' SSP kit. DR4 subtyping was performed using the Dynal DRB1*04-SSP kit (University Transfusion service, Radboud hospital, Nijmegen, The Netherlands).

Peptides

20 Peptides according to the invention and a control peptide were synthesized by solid-phase peptide synthesis. In brief, peptides with free amino- and carboxy termini were synthesized on a Milligen 9050 synthesizer, using Fmoc/tBu protected activated esters on PEG-PS resins. After cleaving off and deprotecting, the peptides were purified by preparative HPLC, converted into acetate salts with Dowex Ac-resin or into chloride salts and lyophilized. The peptides were checked with mass spectrometry. The peptides used in this study are listed in table 1. An N-terminal biotinylated Influenza Haemagglutinine derived peptide (SEQ ID No. 9), biotin-spacer-IHA(307-319)F, in which the third residue (Y) was replaced by F, (biotin-NH-(CH₂)₅-CO-
25 PKFVKQNTLKLAT), was used as marker peptide in the binding studies with DR4Dw4 (DRB1*0401). The non-biotinylated peptide IHA(307-319)F having SEQ ID NO 9 was used as control peptide.

Table 1. The amino acid sequence of the synthesized peptides.

peptide	sequence	HPLC purity	SEQ ID
1	PTFGRSFTLASSE	93.2%	No 5
2	RSFTLASSETGVG	85.6%	No 6
3	VGYYDDQESVKSKV	95.6%	No 7
4	SQRFSKIASNTQSR	97.6%	No 8
IHA(307-319)F	PKFVKQNTLKLAT	97.9%	No 9

The amino acid sequence of peptides 1-4 and the control peptide IHA(307-319)F are given, corresponding to the respective sequence ID's

Cell culture for the production of purified HLA-DR molecules

Two EBV-transformed B-cell lines, BSM (A2, B62, Cw3, DR4Dw4, DQ8, Dpw2) and BM92 (A25, B51, Cw1, DR4Dw14, DQ8), were a gift from the Academic Hospital Leiden, the Netherlands. The cells were cultured in DMEM/HAM's F12 (Gibco Laboratories, Grand Island, NY) supplemented with 10% FCS (Hyclone Laboratories), 1% non-essential amino acids (ICI), L-glutamine, 2-ME and antibiotics. Cells were routinely passaged every 2-3 days in a 1:2 ratio. Cells were harvested and thereafter washed three times in PBS (4°C) containing 1 mM PMSF. Cell pellets were stored at -70°C until use.

Affinity purification of HLA-DR molecules

HLA-DR molecules were affinity purified from cell lysates using monoclonal antibody L243 (ATCC HB55), directed against a nonpolymorphic determinant on the DR-complex (Lampson et al., J. Immunol. 125:293, 1980). Protein G sepharose purified L243 was coupled to NHS-Sepharose 4 FF (Pharmacia) according to the manufactures instructions.

HLA-DR expressing cells were thawed and lysed on ice for 30 minutes in PBS, 1% NP-40, 1 mM AEBSF (Calbiochem). The lysate was cleared by centrifugation at 15 000 rpm (Sorvall, SS34 rotor) for 30 minutes. The supernatant was passed through a 0.45 µm filter and added to L243-NHS-sepharose beads. After overnight incubation, the beads were transferred to a column and washed with five volumes PBS, 1% NP-40, 5 volumes PBS, 0.5% NP-40, 15 volumes PBS, 0.5% NP-40, 0.1% SDS, 5 volumes PBS, 0.05% NP-40, 5 volumes PBS, 1% n-

octyl-glucoside (Sigma, St. Louis, USA) and 5 volumes 50 mM diethylamine (Fluka), 150 mM NaCl, 1% n-octyl-glucoside pH=8.0. HLA-DR molecules were eluted with 50 mM diethylamine, 150 mM NaCl, 1% n-octyl-glucoside pH=11. Immediately after collection, the fractions were neutralised with 2M glycine pH=4.8. Collected fractions were analysed on SDS-PAGE under non-reducing conditions followed by silver staining. Fractions containing purified HLA-DR were concentrated by ultrafiltration over a 30 kD cut-off membrane.

HLA-DR peptide binding assay

The peptide binding studies were performed using an improved version of a semi-quantitative binding assay described previously (Joosten et al., Int. Immunol. 6:751, 1994). Purified HLA-DR molecules (0.5-500 nM) were incubated at pH=5.0 with 50 nM biotinylated marker peptide (biotin-spacer-IHA(307-319)F) and a concentration range of competitor peptide (peptides 1-4 and IHA(307-319)F as a control peptide) in a final volume of 25 µl binding buffer (PBS, 1 mM AEBSF, 1 mM N-ethyl maleimide, 8 mM EDTA, 10 µM pepstatin A, 0.01% NaN₃, 0.05% NP-40 and 5% DMSO).

After approximately 45 hours incubation at room temperature, bound and unbound marker peptides were separated either by SDS-PAGE in combination with blotting on a nitrocellulose filter (BioRad) or by vacuum DOT blotting using a nitrocellulose filter (BioRad) and 96 wells Hybry Dot equipment (BRL). Blots were blocked with 0.5% DNA blocking reagent (Boehringer Mannheim, Germany) in 0.1 M maleic acid pH=7.5, 150 mM NaCl. After 1/2 hour, blots were washed in PBS, 0.02% Tween 20 (Sigma, St. Louis, USA) and incubated with Streptavidin-HRP (Southern Biotechnology) in a 1:40,000 or 1:5,000 dilution respectively. DR-bound, biotinylated marker peptide was detected by enhanced chemoluminescence using a Western Blot ECL kit (Amersham, U.K.) according to the manufactures instructions. Preflashed films (hyperfilm-ECL, Amersham, U.K.) were exposed for 10 minutes. The relative binding affinity of a given peptide was related to competition with the marker peptide. This relative affinity was defined as the peptide concentration at which the signal was reduced to 50% (125 I-IC₅₀).

In case of the SDS-PAGE, 125 I-IC₅₀ values were determined by visual inspection. The density of the DOT Blot-spots was analysed using a computing densitometer (Molecular Dynamics, USA) and Image Quant and Excel software.

Proliferative responses of blood mononuclear cells

In order to identify T cell reactivity to peptide sequences within HC gp-39, the peptides 1-4 were tested for their capacity to induce proliferative responses in PBMC from RA patients versus healthy DR1 or DR4 controls

PBMC obtained from heparinized venous peripheral blood were isolated by standard centrifugation on a Ficoll-Paque gradient. Cells were cultured in three- or four-fold at a concentration of 1.5×10^5 cells/well in medium supplemented with 10% heat-inactivated autologous plasma, L-glutamine, 2-ME and antibiotics in flatbottomed microtiter plates. Cells were incubated in medium alone or in the presence of PHA (2.5 μ g/ml) or in the presence of antigens, including the *Candida albicans* extract (recall antigen) (1 μ g/ml, 0.1 μ g/ml) or one of the peptides 1-4 in concentrations of 100 μ g/ml, 25 μ g/ml or 10 μ g/ml. Cultures were incubated in a total volume of 210 μ l for 7 days at 37 °C in a humidified atmosphere of 5% CO₂. Cultures were pulsed during the last 18 hours with 0.25 μ Ci ³H-thymidine

Definitions

Patients found to respond to both concentrations of at least one peptide tested were ranked as high responders (HR), patients found to respond to at least one of the concentrations of at least one peptide tested were ranked as responders (R). Consequently, patients which did not respond to any of the peptides tested were ranked as non-responders (NR)

The relative binding affinity of a given peptide was related to competition with the marker peptide. This relative affinity was defined as the peptide concentration at which the signal was reduced to 50% (125 I-IC₅₀)

RESULTS

In order to determine T cell reactivity to peptides 1-4, the PBMC proliferative response in RA patients and healthy donors was analyzed

So far, most HR or R to the HC gp-39-derived peptides carried the DR1, DR4Dw4, DR4Dw14 or the DR10 specificity, all of which are known to be associated with an increased risk for the development of RA. Binding of the peptides 1-4 to DR4Dw4 and DR4Dw14 was demonstrated as shown in Table 2

Table 2: Binding of the peptides 1-4 to HLA-DR molecules

Peptide	DR4Dw4		DR4Dw14	
	SDS-PAGE	DOT Blot	SDS-PAGE	DOT Blot
1	0.04	0.04	2.5	1.95
2	0.04	0.08	0.2	0.19
3	1	0.23	≥ 50	> 10
4	-	0.08	-	0.4
IHA(307-319)F	0.3	0.55	0.3	0.53

Binding of peptides 1-4 having amino acid sequence given in SEQ ID No. 5-8 respectively to HLA-DR molecules. Peptide binding was determined in a semiquantitative binding assay (as described in Methods of Example 1) at pH 5, using 50 nM IHA(307-319)F as a marker peptide. The values are IC_{50} values in micromolar (concentration competitor peptide at which the signal is reduced to about 50% of the signal without competitor peptide). All results, except for peptide 4, are from two independent experiments (SDS-PAGE and DOT Blot, respectively).

Most RA patients responded to one or more of the peptides according to the invention (Table 3), whereas responses were rarely found in the healthy donor group (Table 4). 6 RA patients were found not to respond to any of the peptides tested (results not shown).

Disease severity in RA patients was ranked from stage I-IV. The rontgenscore as here indicated is an estimation of the degree of joint destruction. HR and R to the peptides 1-4 were found in all stages of disease.

The peptides according to the invention represent autoreactive T cell epitopes and reactivity to these epitopes was found predominantly in RA patients but rarely in healthy donors. Thus, RA patients have activated, autoreactive T-cells directed at the autoantigen HC gp-39. Clearly the autoantigen HC gp-39 is under attack of T-cells, resulting in inflammation and destruction of the HC gp-39 antigen in the cartilage.

Table 3: Proliferative responses of PBMC from RA patients to peptides 1-4 measured in 10% autologous serum.

Patient	Peptide 1			Peptide 2			Peptide 3			Peptide 4			Disease state
	100 µg/l	25 µg/l	10 µg/l	100 µg/l	25 µg/l	10 µg/l	100 µg/l	25 µg/l	10 µg/l	100 µg/l	25 µg/l	10 µg/l	
215-0 (HR)	13	5		19	3		6	3					III-IV
240-0 (HR)	9	5		6	5		6	2					II-III
242-0 (HR)	-	0		-	4		3	9					I
243-0 (R)	2		<1	6		1	1		1				III
247-0 (R)	9		1	2		1	-		1				III
248-0 (R)	2		1	1		1	1		1				III
191-03 (HR)	3		2	<2		1	<2		1	<2		<2	IV
272-0 (R)	<2		1	<1		<1	<1		<1	6		<2	III-IV
273-0 (HR)	14		12	6		7	6		1	29		3	IV
275-0 (HR)	2		4	9		2	1		1	5		<2	III
276-0 (HR)	4		3	1		3	1		<1	<2		1	IV, burnt out
280-0 (HR)	11		6	4		1	1		1	2		<2	IV, burnt out

Proliferative responses of PBMC from RA patients to the peptides 1-4 with the amino acid sequence given in SI-Q ID No 5-8 respectively.

Results are presented as stimulation index (SI) values (\pm antigen-specific cpm (mean of measurements) control cpm (mean of measurements)). SI values ≥ 2 were regarded as positive. SI values ≥ 2 as negative (\pm standard deviations of the mean of measurements exceeded acceptable values, therefore values are not given). HR = high responder (\pm responses at both peptide concentrations tested); R = responder (\pm responses at the highest peptide concentration tested); Burnt out = almost all cartilage has disappeared.

Table 4. Proliferative responses of freshly isolated PBMC from healthy donors to peptides 1-4 measured in 10 % autologous serum.

Donor	Peptide 1			Peptide 2			Peptide 3			Peptide 4		
	100 µg/ml	10 µg/ml	1 µg/ml	100 µg/ml	10 µg/ml	1 µg/ml	100 µg/ml	10 µg/ml	1 µg/ml	100 µg/ml	10 µg/ml	1 µg/ml
155-03 (NR)	1	1	1	1	1	1	1	<2				
156-03 (NR)	2	2	1	<1	1	<1	<1	1	1	<2	<1	
157-03 (NR)	2	1	1	<2	<2	1	1	1	1			
157-04 (R)	-	10	1	<1	1	-	-	1	1	1	1	
168-02 (NR)	1	<2	<1	<1	<2	<2	<2	<2				
168-03 (NR)	<1	<2	<1	<1	<1	1	1	1	1	<1	1	
169-02 (NR)				1	1	1	1	1				
230-01 (NR)	1	1	1	1	1	1	1	1				
235-0 (NR)	2	1	<2	<2	1	<2	<2	1				
235-01 (NR)	1	<2	<1	<1	<1	<2	<2	<1				
236-01 (NR)	1	<1	1	1	1	<1	<1	1				
237-0 (NR)	1	1	1	<1	<2	1	1	<1				
237-01 (NR)	1	<1	1	1	1	1	1	1	1	<1	<1	
238-0 (NR)	1	1	1	<1	1	1	1	1				
238-01 (NR)	1	<1	1	1	1	1	1	1	1	<1	1	
241-0 (NR)	1	<2	<1	<1	1	<1	<1	<1				
241-01 (R)	2	<1	<1	<1	<1	1	1	<1	<1	<1	<1	

Proliferative responses of PBMC from healthy donors to peptides 1-4 having the amino acid sequences given in SEQ ID NO 5-8 respectively. Results are presented as stimulation index (SI) values (antigen-specific cp5m (mean of measurements) / control cp5m (mean of measurements)). SI values < 2 were regarded as negative. R = responder (+) responses at the highest peptide concentration tested; - standard deviations of the mean of measurements exceeded acceptable values, therefore values are not given.

EXAMPLE 2

METHODS

Purification of HC gp-39 from the MG63 osteosarcoma cell line

MG63 cells (human osteosarcoma ATCC CRL 1427) were cultured in cell factories in DMEM/HAM's F12 serum free medium. HC gp-39 was purified from the culture supernatant by heparin affinity chromatography followed by super dex 75 chromatography. Purity was checked by SDS-PAGE. In addition, N-terminal amino acid sequencing confirmed that the purified protein was identical to the protein described by Hakala et al.

Arthritogenicity of HC gp-39 in Balb/c mice

10 or 50 µg of purified HC gp-39 in a 100 µl volume PBS (0.5 M NaCl, 0.01 M sodium phosphate buffer, pH 7.5) mixed 1:1 in incomplete Freund's adjuvant (IFA) was injected subcutaneously in the chest region in 2 x 4 female Balb/c mice (Harlan CPB, Zeist, The Netherlands) whereas 4 controls were injected with PBS (1:1 in IFA). Mice were examined daily for clinical signs of arthritis. Severity of arthritis was assessed by scoring each paw from 0-3 (according to the article by Glant et al). In short, score 0 = no changes, score 1 = erythema and swelling, score 2 = swelling and appearance of deformities, score 3 = immobility due to loss of flexion and extension.

Tolerance induction by intranasal administration of HC gp-39

Twenty eight µg of protein was administered intranasally (2 x 10 µl) to 10 female Balb/c mice (anesthetized lightly with Enflurane) using a PT45 micro conduit and a Hamilton syringe. Antigen was administered on day -15, -10 and -5 prior to arthritis induction. Controls (n = 10) were submitted to the same procedure but received the vehicle (PBS) only (Table 5).

Immunological tolerance was evaluated by measuring delayed type hypersensitivity (DTH) responses following sensitization as described above on day 0, using 10 µg of protein. Sensitization on day 0 was followed by an injection of 10 µg HC gp-39 in 50 µg volume in the left hind footpad on day 8 (challenge). DTH reactions were measured as the increase in footpad ([swelling left (mm x 10⁻³) - swelling right (mm x 10⁻³)] / swelling right (mm x 10⁻³) x 100%. The

- 20 -

footpad swelling was measured using an inhouse designed μ meter at 0, 24 and 48 hr after challenge

Tolerance to arthritis induction in these same mice was then further monitored until day 31 following sensitization and mice were examined daily for clinical signs. Severity of arthritis was assessed as mentioned above

Table 5. Tolerization scheme.

DAY	HC gp-39 tolerized	non-tolerized
-15	28 μ g HC gp-39 intra nasal	PBS
-10	28 μ g HC gp-39 intra nasal	PBS
-5	28 μ g HC gp-39 intra nasal	PBS
0	10 μ g HC gp-39 subcutaneous	10 μ g HC gp-39 subcutaneous
8	10 μ g HC gp-39 footpad	10 μ g HC gp-39 footpad
9	24 hour DTH	24 hour DTH
10	48 hour DTH	48 hour DTH
0-31	score arthritic signs	score arthritic signs

DTH = delayed type hypersensitivity. PBS = 0.5 M NaCl, 0.01 M sodium phosphate buffer, pH 7.5

RESULTS

Arthritogenicity of HC gp-39

Following one injection of 50 μ g HC gp-39 mixed with IFA, all mice gradually developed a severe arthritis (Table 6). Signs of arthritis were observed first at day 15-20 after sensitization, in the fore paws of 3 out of 4 animals. The mouse that did not show any signs in the fore paws developed arthritis in the hind paws by day 34 upon sensitization. The course of HC gp-39-induced disease was characterized by relapses occurring periodically in fore paws or hind paws and gradually developed from mild arthritis into a more severe arthritis (disease progression was followed for 62 days). Very often (>50%) a symmetrical distribution of afflicted joints was observed, meaning that both fore paws or both hind paws showed arthritic signs at the same time.

- 21 -

Arthritis induction was similarly achieved using 10 μ g instead of 50 μ g HC gp-39 mixed with IFA. Arthritic scores, however, were somewhat lower (data not shown). Three out of 4 mice developed a severe arthritis. One mouse showed only mild signs during the duration of the experiment. All through the length of the experiment control mice showed no signs of arthritis.

In aggregate, both 10 and 50 μ g of protein were sufficient to induce a progressive arthritis in Balb/c mice. The chronic nature of arthritis induction by HC gp-39, characterized by recurrent relapses in addition to symmetrical affliction of joints is reminiscent of disease progression in rheumatoid arthritis (RA).

Table 6. Initiation and progression of arthritis in HC gp-39 sensitized Balb c mice

sensitization (n = 4)	arthritis onset (day)		arthritis signs		
	FP	HP	none	mild	severe
PBS contr.	-	-	4	0	0
10 μ g HC gp-39	-, 13, 15, 57	32, 34, 36, 43	0	1	3
50 μ g HC gp-39	13, 15, 15, 48	29, 34, 34, 53	0	0	4

FP = fore paw HP = hind paw contr = controls. Arthritis signs: none = score = 0, mild = score per animal does not exceed 2, severe = score per animal is \geq 4.

Immunological tolerance measured in the DTH assay

Control mice injected with HC gp-39 at day 0 showed a strong, antigen-specific DTH response, which suggests that a cellular immune response to HC gp-39 was elicited upon sensitization (Table 7). Intranasal administration of HC gp-39, however, completely abrogated DTH responses upon challenge with the autoantigen, thereby showing that the HC gp-39-specific T-cells were indeed tolerized.

Notably, 4 out of 10 animals of the non-tolerized group developed arthritis in the ankle adjacent to the site of challenge. In contrast, the tolerized group did not develop an arthritis in the joints neighbouring the challenged site, thereby suggesting that immunological tolerance to HC gp-39 results in protection against arthritis development.

- 22 -

Table 7. DTH responses to HC gp-39 following tolerization by nasal administration

treatment	mean % swelling			arthritis ankle
	0 hr	24 hr	48 hr	
controls (n = 10)	-1.3	31.4	38.6	4/10
tolerized (n = 9)	-0.05	3.7	1.5	0/9

Tolerance to arthritis induction or progression

The HC gp-39 tolerized and the non-tolerized Balb/c mice were then further monitored for initiation and progression of arthritis

In all mice of the control (non-tolerized) group, disease was initiated upon sensitization with HC gp-39 (Table 8). Seven mice gradually developed a severe arthritis whereas three mice showed only mild signs (highest score 2). In contrast, five mice of the HC gp-39-tolerized group were protected against disease development during the course of the experiment. Furthermore, two animals of the tolerized group showed only mild signs for brief periods of time. Three animals developed a more severe arthritis with scores of 4.

Interestingly, in tolerized animals, arthritis onset was delayed in both hind and fore paws by a minimum of 7-9 days respectively (Figure 1). Although fewer animals were affected in the tolerized group, the arthritic score per animal of the fore paws was comparable to the arthritic score in the non-tolerized animals. The arthritic score per animal of the hind paws however, was somewhat lower in the tolerized animals (Figure 1).

Table 8. Initiation and progression of arthritis in HC gp-39-tolerized and non-tolerized Balb/c mice.

animals	arthritis onset (day)	arthritis signs		
		none	mild	severe
controls (n = 10)	13-16	0	3	7
tolerized (n = 10)	23-24	5	2	3

Onset arthritis: first signs appear - highest number of animals affected. Arthritis signs: no signs: score = 0; mild: score per animal does not exceed 2; severe: score per animal is > 4.

- 23 -

The experiments above demonstrate the arthritogenic nature of HC gp-39 in Balb/c mice. The course of HC gp-39-induced disease was characterized by relapses occurring periodically in fore paws and/or hind paws and gradually developed from a mild arthritis into a more severe form. Also, a symmetrical distribution of afflicted joints was observed which is together with the observation of recurrent relapses, reminiscent of disease progression in rheumatoid arthritis. The fierce arthritogenic nature of HC gp-39 was illustrated by a single, subcutaneous injection of 10 or 50 µg of protein which initiated arthritic signs in all animals. That HC gp-39 specific T cells are indeed elicited in response to HC gp-39 sensitization was shown by induction of HC gp-39-specific DTH responses. These data were further confirmed by the demonstration of HC gp-39-specific *in vitro* proliferative responses in animals immunized in the footpad with HC gp-39 (data not shown). Importantly, non-tolerized animals developed arthritis in the ankle neighbouring the injection site, thereby indeed suggesting an involvement of HC gp-39-specific T cells in arthritis induction.

Intranasal administration of peptide antigen has been used to induce antigen-specific immune tolerance. The experiments showed that intranasal administration of HC gp-39 leads to immunological non-responsiveness. DTH responses following sensitization were completely abrogated in HC gp-39-tolerized mice whereas control mice showed an antigen-specific swelling. These observations indicate that administration of HC gp-39 leads to peripheral immune tolerance.

In non-tolerized animals DTH responses were accompanied by arthritis in the ankle (adjacent to the challenged site) in four out of ten mice. In contrast, the ankles of HC gp-39-tolerized animals were indeed fully protected, thereby suggesting that autoreactive T-cells have been effectively silenced. The notion that tolerization with HC gp-39 protects against disease development was taken further by the observation that 5 out of 10 animals in the tolerized group were entirely protected throughout the length of the experiment. Although the other five animals in the group did eventually develop clinical signs, the onset of arthritis was considerably delayed. Hence it can be concluded that HC gp-39-specific T cells are involved in the arthritogenic process and more importantly, that by tolerization of these T cells with a pharmaceutical composition according to the present invention arthritis development can be delayed or suppressed.

- 24 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: AKZO NOBEL N.V.
(B) STREET: Velperweg 76
(C) CITY: Arnhem
(E) COUNTRY: The Netherlands
(F) POSTAL CODE (ZIP): 6824 BM
(G) TELEPHONE: 0412-666376
(H) TELEFAX: 0412-650592
(I) TELEX: 37503 akpha nl

(ii) TITLE OF INVENTION: Novel Peptides derived from
autoantigen for use in Immunotherapy of Autoimmune Diseases

(iii) NUMBER OF SEQUENCES: 9

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

A LENGTH: 9 amino acids
B TYPE: amino acid
C STRANDEDNESS: single
D TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Phe Gly Arg Ser Phe Thr Leu Ala Ser
1

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

A LENGTH: 9 amino acids
B TYPE: amino acid
C STRANDEDNESS: single
D TOPOLOGY: linear

- 25 -

(11) MOLECULE TYPE: peptide

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Phe Thr Leu Ala Ser Ser Glu Thr Gly
1 5

(2) INFORMATION FOR SEQ ID NO: 3:

(1) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(11) MOLECULE TYPE: peptide

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Tyr Asp Asp Gln Glu Ser Val Lys Ser
1 5

(2) INFORMATION FOR SEQ ID NO: 4:

(1) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(11) MOLECULE TYPE: peptide

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Phe Ser Lys Ile Ala Ser Asn Thr Gln
1 5

(2) INFORMATION FOR SEQ ID NO: 5:

(1) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(11) MOLECULE TYPE: peptide

- 26 -

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Pro Thr Phe Gly Arg Ser Phe Thr Leu Ala Ser Ser Glu
1 5 10

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Arg Ser Phe Thr Leu Ala Ser Ser Glu Thr Gly Val Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

Val Gly Tyr Asp Asp Gln Gln Ser Val Lys Ser Lys Val
1 5 10

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 27 -

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Ser Gln Arg Phe Ser Lys Ile Ala Ser Asn Thr Gln Ser Arg
1 5 10

5

(2) INFORMATION FOR SEQ ID NO: 9:

10

(1) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 13 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15

(11) MOLECULE TYPE: peptide

20

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Pro Lys Phe Val Lys Gln Asn Thr Leu Lys Leu Ala Thr
1 5 10

25

Claims

- 1 Peptide comprising a subsequence of HC gp-39, said peptide comprising one or more of
the amino acid sequences FGRSFTLAS (SEQ ID No 1), FTLASSETG (SEQ ID No 2),
5 YDDQESVKS (SEQ ID No 3) and FSKIASNTQ (SEQ ID No 4)
- 2 Peptide according to claim 1, said peptide comprising one or more of the amino acid
sequences PTFGRSFTLASSE (SEQ ID No 5), RSFTLASSETGVG (SEQ ID No 6),
10 VGYDDQESVKSKV (SEQ ID No 7) and SQRFSKIASNTQSR (SEQ ID No 8)
- 3 Peptide according to claim 1 or 2 having the amino acid sequence PTFGRSFTLASSE
(SEQ ID No 5)
- 4 Peptide according to claim 1 or 2 having the amino acid sequence RSFTLASSETGVG
15 (SEQ ID No 6),
- 5 Peptide according to claim 1 or 2 having amino acid sequence VGYDDQESVKSKV
(SEQ ID No 7)
- 20 6 Peptide according to claim 1 or 2 having amino acid sequence SQRFSKIASNTQSR
(SEQ ID No 8)
- 7 The HC gp-39 protein or peptides according to any of the claims 1 to 6 for use as a
therapeutical substance
- 25 8 Pharmaceutical composition comprising the HC gp-39 protein or one or more of the
peptides according to any of the claims 1 to 6, and a pharmaceutical acceptable carrier
- 9 Use of the HC gp-39 protein or one or more of the peptides according to any of the
30 claims 1 to 6 for the manufacture of a pharmaceutical preparation for the induction of

- 29 -

specific T-cell tolerance to the HC gp-39 autoantigen in patients suffering from T-cell mediated cartilage destruction

10 Use of the HC gp-39 protein or one or more of the peptides according to any of the
5 claims 1 to 6 for the manufacture of a pharmaceutical preparation for the induction of specific T-cell tolerance to the HC gp-39 autoantigen in patients suffering from arthritis, more specifically rheumatoid arthritis

11 HC gp-39 protein or one or more of the peptides according to any of the claims 1-6 for
10 use in a method to induce clinical arthritis in non-human mammals, preferably mice

12 Peptides according to any of the claims 1-6 for use as a diagnostic substance

13 Diagnostic composition comprising one or more of the peptides according to any of the
15 claims 1-6 and a detection agent

14 A diagnostic method for the detection of activated autoreactive T cells comprising the following steps

a) isolation of the peripheral blood mononuclear cells (PBMC) from a blood sample of an
20 individual,

b) culture of said PBMC under suitable conditions,


c) incubation of said PBMC culture in the presence of HC gp-39, fragments thereof and/or one or more peptides according to any of the claims 1-6, and

d) detection of a response of T cells, indicating the presence of activated autoreactive T
25 cells in the individual


15 Test kit for the detection of activated autoreactive T cells, said test kit comprising HC gp-39 or one or more of the peptides according to any of the claims 1-6

30 16 Peptide according to any of the claims 1-15 having an amino acid sequence of 9-55 amino acid residues

1/4

FIG. 1A: Arthritis score fore paw
non tolerized


2/4

FIG. 1B: Arthritis score for paw
tolerized

3/4

FIG. 1C: Arthritis score hind paw


non tolerized



4/4

FIG. 1D: Arthritis score hind paw

tolerized



INTERNATIONAL SEARCH REPORT

International Application No.

PC/EP 95/04201

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07K14/47 A61K38/04 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 268, no. 34, 5 December 1993 BALTIMORE, MD US, pages 25803-25810, B E HAKALA ET AL. 'Human carilage gp-39, a major secretory product of articular chondrocytes and synovial cells, is a mammalian member of chitinase protein family' cited in the application see table 1 --- -/-	1,2,6-10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *A* document member of the same patent family

Date of the actual completion of the international search

19 February 1996

Date of mailing of the international search report

19.03.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Masturzo, P

INTERNATIONAL SEARCH REPORT

International Application No.

PC/EP 95/04201

(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BIOCHEMICAL JOURNAL, vol. 269, no. 1, 1 July 1990 LONDON, THE BIOCHEMICAL SOCIETY, pages 265-268, P NYIRKOS & E E GOLDS 'Human synovial cells secrete a 39 kDa protein similar to a bovine mammary protein expressed during the non-lactating period' see the whole document ---	1-10
X	DATABASE WPI Week 10 1988 Derwent Publications Ltd., London, GB; AN 88-068419 'New polypeptide and DNA coding-related to protective mechanisms such as immune response..' & JP,A,63 023 898 (DAINIPPON PHARM.) , 16 July 1986 see abstract ---	1-10, 12-14
X	DATABASE STRAND Genetics Computer Group Sequence, 1 February 1988 T FURUYA ET AL 'cDNA encoding new polypeptide' see abstract & JP,A,88 023 898 (DAINIPPON PHARM.) 1 February 1988 ---	1-10, 12-14
P,X	WO,A,95 01995 (THE REGENTS OF UNIVERSITY OF CALIFORNIA) 19 January 1995 see the whole document -----	1-16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/EP 95/04201

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9501995	19-01-95	NONE	



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 15/12, C07K 14/47, A61K 38/17, A01K 67/027	A1	(11) International Publication Number: WO 97/40149 (43) International Publication Date: 30 October 1997 (30.10.97)
(21) International Application Number: PCT/EP97/01903 (22) International Filing Date: 15 April 1997 (15.04.97) (30) Priority Data: 08/634,493 18 April 1996 (18.04.96) US (71) Applicant (for all designated States except US): AKZO NOBEL N.V. [NL/NL]; Velperweg 76, NL-6824 BM Arnhem (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): BOOTS, Anna, Maria, Helena [NL/NL]; Verlengde Torenstraat 10, NL-5366 AV Megen (NL). VERHEIJDEN, Gijsbertus, Franciscus, Maria [NL/NL]; Heischouw 7, NL-5345 XT Oss (NL). BOS, Ebo, Sybren [NL/NL]; De Vriesstraat 8, NL-5344 JA Oss (NL). (74) Agent: HERMANS, Franciscus, G., M.; P.O. Box 20, NL-5340 BH Oss (NL).		(81) Designated States: AM, AU, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IS, JP, KG, KP, KR, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: AUTOANTIGEN AND PROTEINS STRUCTURALLY RELATED THERETO FOR USE IN IMMUNOTHERAPY OF AUTOIMMUNE DISEASES		
(57) Abstract The present invention relates to the use of autoantigen HC gp-39, and proteins comprising an amino acid sequence which exhibits at least 50 % homology with the amino acid sequence of HC gp-39, more particular with the amino acid sequence YKLVCYYTSWSQYREGDGSCFPDALDR-FLCTHIIYSFANISND (SEQ ID NO:1) in antigen-specific treatment of articular cartilage destruction in autoimmune diseases in mammals to induce systemic tolerance of the immune system. The autoantigen HC gp-39, and the arthritogenic proteins comprising an amino acid sequence which exhibits at least 50 % homology with the amino acid sequence YKLVCYYTSWSQYREGDGSCFPDALDR-FLCTHIIYSFANISND (SEQ ID NO:1) are also suitable to induce arthritis in animals, preferably mice. The invention furthermore relates to pharmaceutical compositions comprising said autoantigen and/or said arthritogenic proteins, a diagnostic method for the detection of autoreactive T cells in a test sample and test kits to be used in said method.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

AUTOANTIGEN AND PROTEINS STRUCTURALLY RELATED THERETO FOR USE IN IMMUNOTHERAPY OF AUTOIMMUNE DISEASES

5

The invention relates to a novel autoantigen and proteins related thereto, their use in treatment of chronic destruction of articular cartilage in autoimmune diseases, pharmaceutical compositions comprising said autoantigen and/or proteins, a diagnostic method for the detection of autoreactive T cells in a test sample and test kits to be used in said method.

10

The immune system is established on a principle of discrimination between foreign antigens (non-self antigens) and autoantigens (self antigens, derived from the individuals own body) achieved by a build in tolerance against the autoantigens.

The immune system protects individuals against foreign antigens and responds to exposure to a foreign antigen by activating specific cells such as T- and B lymphocytes and producing soluble factors like interleukins, antibodies and complement factors. The antigen to which the immune system responds is degraded by the antigen presenting cells (APCs) and a fragment of the antigen is expressed on the cell surface associated with a major histocompatibility complex (MHC) class II glycoprotein. The MHC-glycoprotein-antigen-fragment complex is presented to a T cell which by virtue of its T cell receptor recognizes the antigen fragment conjointly with the MHC class II protein to which it is bound. The T cell becomes activated, i.e. proliferates and/or produces interleukins, resulting in the expansion of the activated lymphocytes directed to the antigen under attack (Grey et al., Sci. Am., 261:38-46, 1989).

Self antigens are also continuously processed and presented as antigen fragments by the MHC glycoproteins to T cells (Jardetsky et al., Nature 353:326-329, 1991). Self recognition thus is intrinsic to the immune system. Under normal circumstances the immune system is tolerant to self antigens and activation of the immune response by these self antigens is avoided.

When tolerance to self antigens is lost, the immune system becomes activated against one or more self antigens, resulting in the activation of autoreactive T cells and the production of autoantibodies. This phenomenon is referred to as autoimmunity. As the immune response in

general is destructive, i.e. meant to destroy the invasive foreign antigen, autoimmune responses can cause destruction of the body's own tissue.

5 The contribution of T cells to autoimmune diseases has been established by several studies. In mice, experimental autoimmune encephalomyelitis (EAE) is mediated by a highly restricted group of T cells, linked by their specificity for a single epitope of myelin basic protein (MBP) complexed to an MHC class II molecule. In the Lewis rat, a species with high susceptibility to various autoimmune diseases, disease has been shown to be mediated by T cells. In humans autoimmune diseases are also thought to be associated with the development of
10 auto-aggressive T cells.

A destructive autoimmune response has been implicated in various diseases such as rheumatoid arthritis (RA), in which the integrity of articular cartilage is destroyed by a chronic inflammatory process resulting from the presence of large numbers of activated lymphocytes
15 and MHC class II expressing cells. The mere presence of cartilage appears necessary for sustaining the local inflammatory response: it has been shown that cartilage degradation is associated with the activity of cartilage-responsive autoreactive T cells in RA (Sigall et al., Clin. Exp. Rheumat. 6:59, 1988; Glant et al., Biochem. Soc. Trans. 18:796, 1990; Burmester et al., Rheumatoid arthritis Smolen, Kalden, Maini (Eds) Springer-Verlag Berlin Heidelberg,
20 1992). Furthermore, removal of cartilage from RA patients by surgery was shown to reduce the inflammatory process (G.S. Panayi et al, Clin. Exp. Rheumatol. 11(suppl.8): S1-S8, 1993). The cartilage proteins are therefore considered to be target autoantigens which are competent of stimulating T cells. Activation of these autoreactive T cells leads to development of autoimmune disease. However, the identification of the autoantigenic components that play a
25 role in the onset of rheumatoid arthritis has so far remained elusory.

The inflammatory response resulting in the destruction of the cartilage can be treated by several drugs, such as for example steroid drugs. However, these drugs are often immunosuppressive drugs that are nonspecific and have toxic side effects. The disadvantages of
30 nonspecific immunosuppression makes this a highly unfavourable therapy.

The antigen-specific, nontoxic immunosuppression therapy provides a very attractive alternative for the nonspecific immunosuppression. This antigen-specific therapy involves the treatment of patients with the target autoantigen or antigens having an amino acid sequence which exhibits sequence homology with the amino acid sequence of the target autoantigen, or synthetic T cell-reactive peptides derived from said autoantigen or said antigen. These synthetic peptides correspond to T cell epitopes of the autoantigen and can be used to induce specific T cell tolerance both to themselves and to the autoantigen. Although it seems paradoxical to desensitize the immune system with the very same antigen responsible for activating the immune system, the controlled administration of the target (auto)antigen can be very effective in desensitization of the immune system. Desensitization or immunological tolerance of the immune system is based on the long-observed phenomenon that animals which have been fed or have inhaled an antigen or epitope are less capable of developing a systemic immune response towards said antigen or epitope when said antigen or epitope is introduced via a systemic route.

To effectively use the tolerance therapy to treat the T cell mediated cartilage destruction, there is a great need to identify the responsible autoantigen or antigens having an amino acid sequence which exhibits sequence homology with the amino acid sequence of the target autoantigen, to desensitize patients against the autoantigen that is activating the T cells responsible for the inflammatory process.

It is an object of the invention to provide the autoantigen, and proteins having an amino acid sequence which exhibit sufficient homology with the amino acid sequence of the autoantigen, said autoantigen and/or protein being suitable to induce specific T cell tolerance to the responsible cartilage antigen in patients suffering from T cell-mediated cartilage destruction. It is a further object of the invention to provide arthritic animal testmodels which are suitable for use in screening for novel drugs to suppress arthritic symptoms. It is another object of the invention to provide a method for detecting autoreactive T cells involved in the destruction of articular cartilage and test kits to be used in said method.

It was surprisingly found that Human Cartilage glycoprotein 39 (herein after referred to as HC gp-39) is a target autoantigen in RA patients which activates specific T cells, thus causing or mediating the inflammatory process. The arthritogenic nature of HC gp-39 was

substantiated in the Balb/c mouse. A single, subcutaneous injection of said protein in Balb/c mice was able to initiate arthritic signs in the animals. The course of the HC gp-39- induced disease was characterized by relapses occurring periodically in fore paws and/or hind paws and gradually developed from a mild arthritis into a more severe form. Also, a symmetrical distribution of afflicted joints was observed which is, together with the observation of recurrent relapses and nodule formation, reminiscent of disease progression in arthritis, especially RA.

Even more surprisingly it was found that administration of HC gp-39 resulted in immunological tolerance and, more importantly, in delayed and/or suppressed arthritic development.

It was furthermore surprisingly found that proteins comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39 are able to induce arthritis when injected to animals. In particular proteins comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence YKLVCYYTSWSQYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID NO: 1) were able to induce arthritis in animals, in the same way as described for HC gp-39.

Preferably, these arthritogenic proteins comprise an amino acid sequence which exhibits at least 70%, more preferably 80%, most preferably 90% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTSWSQYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID NO: 1).

The percentage of sequence homology of the arthritogenic proteins according to the invention with the amino acid sequence of SEQ ID NO: 1 is understood to be measured via generally used sequence comparison programmes such as FASTA (W.R. Pearson and D.J. Lipman, Proc. Natl. Acad. Sci. USA, 85, 2444-2448, 1988).

Suitable arthritogenic proteins according to the invention are for example pig heparine-binding 38kDa protein, bovine 39 kDa whey protein, human YKL-39 protein, murine breast regressing 39kDa protein (brp39), human oviduct-specific glycoprotein, murine oviduct-specific glycoprotein, hamster oviduct-specific glycoprotein, bovine oviduct-specific glycoprotein, human chitotriosidase precursor protein and murine secretory protein YM-1 precursor. The arthritogenic proteins according to the invention are very suitable for inducing systemic tolerance of the immune system to homologous autoantigens and can be used to delay and/or suppress arthritic development in mammals.

HC gp-39 is present in serum of both patients and healthy adults, although the serum concentration of the protein is about twice as much in patients as compared to healthy adults. Furthermore, mRNA coding for HC gp-39 can be found in synovial specimens or cartilage obtained from RA patients, whereas cartilage of healthy adults, obtained at surgery, does not contain a significant amount of said mRNA. When articular chondrocytes and synovial cells are cultured, their major secretory product becomes HC gp-39 (Hakala et al., J. Biol. Chem., Vol. 268, 34:25803, 1993). The arthritogenic nature of HC gp-39 was neither described nor suggested in the Hakala et al publication, nor in any other publication.

Proteins of which the amino acid sequence exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTWSQYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID NO: 1) have been described. The identification of pig heparine-binding 38kDa protein is described in Shackelton et al. (1995), J.Biol.Chem. Vol. 270, No. 22, 13076-13083, however no function of the protein was identified. The isolation and characterization of bovine 39 kDa whey protein is described in J.J. Rejman et al. (1988), Biochem.Biophys.Res.Comm. Vol. 150, No. 1, 329-334. Murine breast regressing 39kDa protein (brp39) is described in Morrison and Leder, 1994, Oncogene 9, 3417. Cloning of the cDNA encoding human oviduct-specific glycoprotein and the corresponding amino acid sequence is described in Arias et al. (1994), Biology of Reproduction 51, 685-694. Other mammalian oviduct-specific glycoproteins such as murine- and hamster oviduct-specific glycoprotein are disclosed in JP-A-07107979, Kinosei Peptide Kenkyusho KK. The purification and molecular cloning of the bovine oviduct-specific glycoprotein is described in Y. Sendai et al, 1994, Biol. of Reprod. 50, 927-934. Human chitotriosidase precursor protein is secreted by activated human macrophages and the cloning of the corresponding cDNA and amino acid sequence is described in Boot et al. (1995), J.Biol.Chem. Vol. 270, No. 44, 26252-26256. The amino acid sequence of murine secretory protein YM-1 precursor was submitted to EMBL Data Library, June 1992, Accession No. M94584 by Chang et al., unpublished. The amino acid sequence of human chondrocyte protein YKL-39 is described in Hu et al. (1996), J.Biol.Chem. Vol 271, No. 32, 19415-19420. None of the afore-mentioned publications however hint or suggest towards the arthritogenic nature of the proteins according to the invention nor to the fact that these proteins can be used as a medical substance in a therapy to

induce specific T-cell tolerance to HC gp-39 in mammals, more specifically man, suffering from T-cell mediated cartilage destruction, such as for example arthritis, more specifically rheumatoid arthritis.

5 Thus, according to the invention HC gp-39, and proteins comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTSWSQYREGDGSCFPDAL-
10 DRFLCTHIIYSFANISND (SEQ ID NO: 1) are very suitable for use in a therapy to induce specific T-cell tolerance to HC gp-39 in patients suffering from T-cell mediated cartilage destruction, such as for example arthritis, more specifically rheumatoid arthritis. Also within the scope of the invention are fragments of HC gp-39 and the arthritogenic proteins according to the invention, which are able to induce T-cell specific tolerance to the HC gp-39 autoantigen in the cartilage under attack.

15 WO 95/01995 and WO 95/02188 describe the diagnostic use of HC gp-39 as a marker for RA, however the arthritogenic nature of HC gp-39 is neither disclosed nor suggested. Nowhere do they hint or suggest towards the use of HC gp-39, or fragments thereof or T-cell reactive peptides according to the present invention in the antigen specific therapy to induce T-cell specific tolerance to the HC gp-39 in the cartilage under attack.

20 HC gp-39, arthritogenic proteins comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTSWSQYREGDGSC-FPDALDRFLCTHIIYSFANISND (SEQ ID NO: 1), and fragments derived from said arthritogenic proteins according to the
25 invention can be prepared with the aid of recombinant DNA techniques. For this purpose, a nucleic acid sequence which codes for HC gp-39 or a protein comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTSWSQYREGDGSCFPDALDR-
30 FLCTHIIYSFANISND (SEQ ID NO: 1) or a fragment derived from the proteins according to the invention or a multimer of said fragment is inserted into an expression vector. Suitable expression vectors are, amongst others, plasmids, cosmids, viruses and YAC's (Yeast Artificial

Chromosomes) which comprise the necessary control regions for replication and expression. The expression vector can be brought to expression in a host cell. Suitable host cells are, for instance, bacteria, yeast cells and mammalian cells. Such techniques are well known in the art (Sambrook et al., Molecular Cloning: a Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, 1989).

Using these recombinant DNA techniques bovine whey protein, the sequence of which has been disclosed herein, can be prepared in a host cell.

Thus, the present invention also provides for isolated cDNA encoding the bovine whey protein as indicated by SEQ ID NO: 3 as well as the bovine whey protein according to SEQ ID NO: 2. It will be clear that also fragments can be prepared once given the teachings of the present invention. The term "fragment" refers to any sequence of amino acids that is part of the protein, having common elements of origin, structure and mechanism of action that are within the scope of the present invention.

Suitable fragments derived from the arthritogenic proteins according to the invention can also be prepared by means of one of the known organic chemical methods for peptide synthesis. The organic chemical methods for peptide synthesis are considered to include the coupling of the required amino acids by means of a condensation reaction, either in homogeneous phase or with the aid of a so-called solid phase. The most common methods for the above condensation reactions are: the carbodiimide method, the azide method, the mixed anhydride method and the method using activated esters, such as described in The Peptides, Analysis, Synthesis, Biology Vol. 1-3 (Ed. Gross, E. and Meienhofer, J.) 1979, 1980, 1981 (Academic Press, Inc.).

Preparation of suitable fragments of above-mentioned HC gp-39 and arthritogenic proteins according to the invention using the "solid phase" is for instance described in J. Amer. Chem. Soc. 85:2149 (1963) and Int. J. Peptide Protein Res. 35:161-214 (1990).

A particular suitable solid phase is, for example, the p-alkoxybenzyl alcohol resin (4-hydroxy-methyl-phenoxy-methyl-copolystyrene-1% divinylbenzene resin), described by Wang (1974) J. Am. Chem. Soc. 95:1328. After synthesis the peptides can be split from this solid phase under mild conditions.

After synthesis of the desired amino acid sequence, detaching of the peptide from the resin follows, for example, with trifluoroacetic acid, containing scavengers, for example triisopropyl silane, anisole or ethanedithiol, thioanisole.

The reactive groups which may not participate in the condensation reaction are, as stated
5 effectively protected by groups which can be removed agains very easily by hydrolisis with the aid of acid, base or reduction. A more extensive account of possible protecting groups can be found in The Peptides, Analysis, Synthesis, Biology, Vol. 1 - 9 (Eds. Gross, Udenfriend and Meienhofer) 1979 - 1987 (Academic Press, Inc.). The protective groups can be split off by
10 various conventional methods, depending on the nature of the particular group, for example with the aid of trifluoroacetic acid or by mild reduction, for example with hydrogen and a catalyst, such as palladium, or with HBr in glacial acetic acid.

Although it seems paradoxical to desensitize the immune system with the very same
15 antigen responsible for activating the immune system, this desensitization is based on the long-observed phenomenon that animals which have been fed or have inhaled an antigen or epitope are less capable of developping a systemic immune response towards said antigen or epitope when said antigen or epitope is introduced via a systemic route. Hence, controlled administration of HC gp-39, and/or proteins comprising an amino acid sequence which exhibits
20 at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTWSQYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID NO: 1) can be effective in desensitization of the immune system. Fragments of the proteins according to the invention that are capable to desensitize patients against HC gp-39 are also within the scope of the invention.

25 According to the invention, patients in which the cartilage is under attack of autoreponsive T cells can be treated with a pharmaceutical composition comprising HC gp-39, one or more proteins comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTWSQYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID NO: 1) or one or
30 more fragments derived from a protein according to the invention and a pharmaceutical

acceptable carrier in order to make the specific autoreactive T cells of these patients tolerant to the HC gp-39 in the cartilage under attack and to diminish the inflammatory response.

Suitable proteins to be used in a pharmaceutical composition according to the invention are proteins comprising an amino acid sequence which exhibits at least 70%, preferably 80%,
5 more preferably 90% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTSWSQYREGDGSCFPDALDRFLCT-HIIYSFANISND (SEQ ID NO: 1).

Very suitable proteins to be used in a pharmaceutical composition according to the invention are for example pig heparine-binding 38kDa protein, bovine 39 kDa whey protein,
10 murine breast regressing 39 kDa protein (brp39), murine oviduct-specific glycoprotein, hamster oviduct-specific glycoprotein, bovine oviduct-specific glycoprotein, human YKL-39, human oviduct-specific glycoprotein, human chitotriosidase precursors protein and murine secretory proteinYM-1 precursor.

Also very suitable to be used in a pharmaceutical composition according to the
15 invention are DNA (expression)vectors comprising DNA which encodes for HC gp-39 or a protein comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTSWS-QYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID NO: 1) or one or more fragments derived from the proteins according to the invention. Upon delivery the
20 DNA (expression)vector can provide by expression a level of the recombinant HC gp-39 or protein comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTSWSQYREGDGSCFP-DALDRFLCTHIIYSFANISND (SEQ ID NO: 1) or fragments according to the invention which is similar to the level which would be achieved by
25 direct administration of a pharmaceutical composition comprising the HC gp-39 protein or peptides.

The autoantigen HC gp-39, arthritogenic proteins comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTSWSQYREGDGSCFPDALDR-
30 FLCTHIIYSFANISND (SEQ ID NO: 1) and fragments derived from said arthritogenic proteins according to the invention have the advantage that they have a specific tolerizing effect on the

autoreactive T cells thus leaving the other components of the immune system intact as compared to the nonspecific suppressive effect of the immunosuppressive steroid drugs. Treatment with the autoantigen or the proteins comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular
5 with the amino acid sequence YKLVCYYTSWSQYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID NO: 1) or fragments according to the invention will be safe and no toxic side effects will occur.

Tolerance can be attained by administering high or low doses of the autoantigen or proteins according to the invention. The amount of autoantigen or protein will depend on the
10 route of administration, the time of administration, the age of the patient as well as general health conditions and diet.

In general, a dosage of 0.01 to 1000 µg of protein per kg body weight, preferably 0.5 to 500 µg, more preferably 0.1 to 100 µg of protein can be used.

Pharmaceutical acceptable carriers are well known to those skilled in the art and include,
15 for example, sterile saline, lactose, sucrose, calcium phosphate, gelatin, dextrin, agar, pectin, peanut oil, olive oil, sesame oil and water. Other carriers may be, for example MHC class II molecules, if desired embedded in liposomes.

In addition the pharmaceutical composition according to the invention may comprise one or more adjuvants. Suitable adjuvants include, amongst others, aluminium hydroxide, aluminium phosphate, amphigen, tocophenols, monophosphoryl lipid A, muramyl dipeptide
20 and saponins such as Quil A. The amount of adjuvant depends on the nature of the adjuvant itself.

Furthermore the pharmaceutical composition according to the invention may comprise one or more stabilizers such as, for example, carbohydrates including sorbitol, mannitol, starch, sucrodedextrin and glucose, proteins such as albumin or casein, and buffers like alkaline
25 phosphates.

Suitable administration routes are intramuscular injections, subcutaneous injections, intravenous injections or intraperitoneal injections, oral and intranasal administration. Oral and intranasal administration are preferred administration routes.

Due to its arthritogenic nature, HC gp-39 as well as the proteins comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTWSQYREGDGSCFPDALDR-FLCTHIIYSFANISND (SEQ ID NO: 1) can be used to induce clinical arthritis in animals. Upon administration of small amounts of HC gp-39 or one or more proteins comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTWSQYREGD-GSCFPDALDRFLCTHIIYSFANISND (SEQ ID NO: 1) or one or more fragments of the proteins according to the invention, arthritic signs will develop in said animals resulting in a disease pattern reminiscent of disease progression in arthritis, especially rheumatoid arthritis. When Balb/c mice were injected subcutaneously with HC gp-39 protein or a protein comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTWSQYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID NO: 1), the animals developed arthritic signs. The course of the HC gp-39-induced disease as well as the disease induced by the protein comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTWSQYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID NO: 1) was characterized by relapses occurring periodically in fore paws and/or hind paws and the gradually development from a mild arthritis into a more severe form. Also, a symmetrical distribution of afflicted joints was observed which is, together with the observation of recurrent relapses and nodule formation, reminiscent of disease progression in arthritis, especially RA.

Thus, these afflicted animals provide an adequate animal model to study the mechanism underlying the initiation and progression of arthritic development. Additionally, said afflicted animals can be used to search for new drugs to treat arthritis and to study the effect of these drugs upon the arthritic development. Preferably mice are used as animal model for arthritis, especially rheumatoid arthritis.

To induce arthritis in said animals, suitable amounts of HC gp-39 or one or more of the proteins according to the invention have to be administered. Suitable amounts are 0.1-1000 µg, preferably 1-100 µg, more preferably 10-50 µg per kg body weight. The amount of HC gp-39 or proteins comprising an amino acid sequence which exhibits at least 50% homology with the

amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTWSQYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID NO: 1) or fragments thereof will depend on the route of administration, time of administration and the type of animal used. Suitable administration routes are the same as described before. To induce the effect of arthritis induction, the HC gp-39 protein, or proteins comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTWSQYREGDGSCFPDALDRFLC-THIIYSFANISND (SEQ ID NO: 1) or peptides according to the invention may comprise one or more stabilizers or adjuvants as described before.

HC gp-39, the proteins comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTWSQYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID NO: 1) or fragments thereof according to the invention are also very suitable for use in a diagnostic method to detect the presence of activated autoreactive T cells involved in the chronic inflammation of the articular cartilage.

The diagnostic method according to the invention comprises the following steps:

- a) isolation of the peripheral blood mononuclear cells (PBMC) from a blood sample of an individual,
- b) culture said PBMC under suitable conditions,
- c) incubation of said PBMC culture in the presence of the autoantigen or proteins according to the invention and fragments thereof, and
- d) detection of a response of T cells, for example a proliferative response, indicating the presence of activated autoreactive T cells in the individual.

In case of detection of a response by measuring the proliferative response of the autoreactive T cells, the incorporation of a radioisotope such as for example ^3H -thymidine is a measure for the proliferation. A response of the autoreactive T cells present in the PBMC can also be detected by measuring the cytokine release with cytokine-specific ELISA, or the cytotoxicity with ^{51}Cr Chromium release. Another detection method is the measurement of expression of activation markers by FACS analysis, for example of IL-2R. A diagnostic

composition comprising one or more of the peptides according to the invention and a suitable detecting agent thus forms part of the invention. Depending on the type of detection, the detection agent can be a radioisotope, an enzyme, or antibodies specific for cell surface or activation markers.

5 Also within the scope of the invention are test kits which comprise one or more peptides according to the invention. These test kits are suitable for use in a diagnostic method according to the invention.

Thus, the present invention provides for a method to detect whether autoaggressive T cells reactive towards HC gp-39 are present in patients suffering from T-cell mediated cartilage
10 destruction such as for example arthritis, in particular rheumatoid arthritis. If HC gp-39-specific T cells are present, tolerization of these T cells with a pharmaceutical composition comprising HC gp-39 or one or more arthritogenic proteins comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence of HC gp-39, more in particular with the amino acid sequence YKLVCYYTWSQYREGDGSCFPDALDRFLCTHIIYS-
15 FANISND (SEQ ID NO: 1) or peptides according to the present invention or combinations thereof can delay or suppress arthritis development.

The following examples are illustrative for the invention and should in no way be interpreted as limiting the scope of the invention.

20 LEGENDS TO THE FIGURES

Figure 1: Initiation and progression of arthritis in HC gp-39 tolerized and non-tolerized Balb/c mice. AS = total arthritic score of the afflicted animals per day following sensitization. N
25 = number of afflicted animals per day following sensitization.

Figure 2: Initiation and progression of arthritis in Balb/c mice upon administration of 0.2 µg, 1 µg, 5 µg and 25 µg bovine 39 kDa whey protein respectively. Initiation/progression is presented as accumulative score (total arthritic score of afflicted animals) per day following
30 sensitization.

EXAMPLE 1

METHODS

Purification of HC gp-39 from the MG63 osteosarcoma cell line

5 MG63 cells (human osteosarcoma ATCC CRL 1427) were cultured in cell factories in DMEM/HAM's F12 serum free medium. HC gp-39 was purified from the culture supernatant by heparin affinity chromatography followed by super dex 75 chromatography. Purity was checked by SDS-PAGE. In addition, N-terminal amino acid sequencing confirmed that the purified protein was identical to the protein described by Hakala et al.

Arthritogenicity of HC gp-39 in Balb/c mice

10 10 or 50 µg of purified HC gp-39 in a 100 µl volume PBS (0.5 M NaCl, 0.01 M sodium phosphate buffer, pH 7.5) mixed 1:1 in incomplete Freund's adjuvant (IFA) was injected subcutaneously in the chest region in 2 x 4 female Balb/c mice (Harlan CPB, Zeist, The Netherlands) whereas 4 controls were injected with PBS (1:1 in IFA). Mice were examined daily for clinical signs of arthritis. Severity of arthritis was assessed by scoring each paw from 0-3 (according to the article by Glant et al). In short, score 0 = no changes, score 1 = erythema and swelling, score 2 = swelling and appearance of deformities, score 3 = immobility due to loss of flexion and extension.

Tolerance induction by intranasal administration of HC gp-39

20 Twenty eight µg of protein was administered intranasally (2 x 10 µl) to 10 female Balb/c mice (anesthetized lightly with Enflurane) using a PT45 micro conduit and a Hamilton syringe. Antigen was administered on day -15, -10 and -5 prior to arthritis induction. Controls (n = 10) were submitted to the same procedure but received the vehicle (PBS) only (Table 1).

Immunological tolerance was evaluated by measuring delayed type hypersensitivity (DTH) responses following sensitization as described above on day 0, using 10 µg of protein. Sensitization on day 0 was followed by an injection of 10 µg HC gp-39 in 50 µg volume in the left hind footpad on day 8 (challenge). DTH reactions were measured as the increase in footpad 30
$$([\text{swelling left (mm} \times 10^{-3}) - \text{swelling right (mm} \times 10^{-3})] / \text{swelling right (mm} \times 10^{-3})) \times 100\%.$$

The footpad swelling was measured using an inhouse designed μ meter at 0, 24 and 48 hr after challenge.

Tolerance to arthritis induction in these same mice was then further monitored until day 31 following sensitization and mice were examined daily for clinical signs. Severity of arthritis was assessed as mentioned above.

Table 1. Tolerization scheme.

DAY	HC gp-39 tolerized	non-tolerized
-15	28 μ g HC gp-39 intra nasal	PBS
-10	28 μ g HC gp-39 intra nasal	PBS
-5	28 μ g HC gp-39 intra nasal	PBS
0	10 μ g HC gp-39 subcutaneous	10 μ g HC gp-39 subcutaneous
8	10 μ g HC gp-39 footpad	10 μ g HC gp-39 footpad
9	24 hour DTH	24 hour DTH
10	48 hour DTH	48 hour DTH
0-31	score arthritic signs	score arthritic signs

DTH = delayed type hypersensitivity. PBS = 0.5 M NaCl, 0.01 M sodium phosphate buffer, pH 7.5

RESULTS

Arthritogenicity of HC gp-39

Following one injection of 50 μ g HC gp-39 mixed with IFA, all mice gradually developed a severe arthritis (Table 2). Signs of arthritis were observed first at day 15-20 after sensitization, in the fore paws of 3 out of 4 animals. The mouse that did not show any signs in the fore paws developed arthritis in the hind paws by day 34 upon sensitization. The course of HC gp-39-induced disease was characterized by relapses occurring periodically in fore paws or hind paws and gradually developed from mild arthritis into a more severe arthritis (disease progression was followed for 62 days). Very often (>50%) a symmetrical distribution of afflicted joints was observed, meaning that both fore paws or both hind paws showed arthritic signs at the same time.

Arthritis induction was similarly achieved using 10 µg instead of 50 µg HC gp-39 mixed with IFA. Arthritic scores, however, were somewhat lower (data not shown). Three out of 4 mice developed a severe arthritis. One mouse showed only mild signs during the duration of the experiment. All through the length of the experiment control mice showed no signs of arthritis.

In aggregate, both 10 and 50 µg of protein were sufficient to induce a progressive arthritis in Balb/c mice. The chronic nature of arthritis induction by HC gp-39, characterized by recurrent relapses in addition to symmetrical affliction of joints is reminiscent of disease progression in rheumatoid arthritis (RA).

Table 2. Initiation and progression of arthritis in HC gp-39 sensitized Balb/c mice.

sensitization (n = 4)	arthritis onset (day)		arthritis signs		
	FP	HP	none	mild	severe
PBS contr.	-	-	4	0	0
10 µg HC gp-39	-, 13, 15, 57	32, 34, 36, 43	0	1	3
50 µg HC gp-39	13, 15, 15, 48	29, 34, 34, 53	0	0	4

FP = fore paw. HP = hind paw. contr. = controls. Arthritis signs: none: score = 0, mild: score per animal does not exceed 2, severe: score per animal is ≥ 3 .

Immunological tolerance measured in the DTH assay

Control mice injected with HC gp-39 at day 0 showed a strong, antigen-specific DTH response, which suggests that a cellular immune response to HC gp-39 was elicited upon sensitization (Table 3). Intranasal administration of HC gp-39, however, completely abrogated DTH responses upon challenge with the autoantigen, thereby showing that the HC gp-39-specific T-cells were indeed tolerized.

Notably, 4 out of 10 animals of the non-tolerized group developed arthritis in the ankle adjacent to the site of challenge. In contrast, the tolerized group did not develop an arthritis in the joints neighbouring the challenged site, thereby suggesting that immunological tolerance to HC gp-39 results in protection against arthritis development.

Table 3. DTH responses to HC gp-39 following tolerization by nasal administration.

teatment	mean % swelling			arthritis ankle
	0 hr	24 hr	48 hr	
controls (n = 10)	-1.3	31.4	38.6	4/10
tolerized (n = 9)	-0.05	3.7	1.5	0/9

Tolerance to arthritis induction or progression

The HC gp-39 tolerized and the non-tolerized Balb/c mice were then further monitored for initiation and progression of arthritis.

In all mice of the control (non-tolerized) group, disease was initiated upon sensitization with HC gp-39 (Table 4). Seven mice gradually developed a severe arthritis whereas three mice showed only mild signs (highest score 2). In contrast, five mice of the HC gp-39-tolerized group were protected against disease development during the course of the experiment. Furthermore, two animals of the tolerized group showed only mild signs for brief periods of time. Three animals developed a more severe arthritis with scores of 4.

Interestingly, in tolerized animals, arthritis onset was delayed in both hind and fore paws by a minimum of 7-9 days respectively (Figure 1). Although fewer animals were affected in the tolerized group, the arthritic score per animal of the fore paws was comparable to the arthritic score in the non-tolerized animals. The arthritic score per animal of the hind paws however, was somewhat lower in the tolerized animals (Figure 1).

Table 4. Initiation and progression of arthritis in HC gp-39-tolerized and non-tolerized Balb/c mice.

animals	arthritis onset (day)	arthritis signs		
		none	mild	severe
controls (n = 10)	13-16	0	3	7
tolerized (n = 10)	23-24	5	2	3

Onset arthritis: first signs appear - highest number of animals affected. Arthritis signs: no signs: score = 0, mild score per animal does not exceed 2, severe: score per animal is ≥ 4

5 The experiments above demonstrate the arthritogenic nature of HC gp-39 in Balb/c mice. The course of HC gp-39-induced disease was characterized by relapses occurring periodically in fore paws and/or hind paws and gradually developed from a mild arthritis into a more severe form. Also, a symmetrical distribution of afflicted joints was observed which is together with the observation of recurrent relapses, reminiscent of disease progression in rheumatoid arthritis. The fierce arthritogenic nature of HC gp-39 was illustrated by a single, subcutaneous injection of 10 or 50 µg of protein which initiated arthritic signs in all animals. That HC gp-39 specific T cells are indeed elicited in response to HC gp-39 sensitization was shown by induction of HC gp-39-specific DTH responses. These data were further confirmed by the demonstration of HC gp-39-specific in vitro proliferative responses in animals immunized in the footpad with HC gp-39 (data not shown). Importantly, non-tolerized animals developed arthritis in the ankle neighbouring the injection site, thereby indeed suggesting an involvement of HC gp-39-specific T cells in arthritis induction.

15 Intranasal administration of peptide antigen has been used to induce antigen-specific immune tolerance. The experiments showed that intranasal administration of HC gp-39 leads to immunological non-responsiveness. DTH responses following sensitization were completely abrogated in HC gp-39-tolerized mice whereas control mice showed an antigen-specific swelling. These observations indicate that administration of HC gp-39 leads to peripheral immune tolerance.

20 In non-tolerized animals DTH responses were accompanied by arthritis in the ankle (adjacent to the challenged site) in four out of ten mice. In contrast, the ankles of HC gp-39-tolerized animals were indeed fully protected, thereby suggesting that autoreactive T-cells have been effectively silenced. The notion that tolerization with HC gp-39 protects against disease development was taken further by the observation that 5 out of 10 animals in the tolerized group were entirely protected throughout the length of the experiment. Although the other five animals in the group did eventually develop clinical signs, the onset of arthritis was considerably delayed. Hence it can be concluded that HC gp-39-specific T cells are involved in the arthritogenic process and more importantly, that by tolerization of these T cells with a pharmaceutical composition according to the present invention arthritis development can be delayed or suppressed.

EXAMPLE 2**METHODS****Arthritogenicity of Bovine 39 kDa whey protein in Balb/c mice**

0.2, 1, 5 or 25 µg of purified Bovine 39 kDa whey protein in a 100 µl volume (0.5 M NaCl, 0.01M sodium phosphate buffer, pH 7.5) mixed 1:1 in incomplete Freund's adjuvant (IFA) was injected subcutaneously in the chest region in 4 x 10 female Balb/c mice (Charles River, Sulzfeld, Germany) whereas 10 controls were injected with PBS (1:1 in IFA). Mice were examined every other day for clinical signs of arthritis. Severity of arthritis was assessed by scoring each paw from 0-3 (according to the article by Glant et al). In short, score 0 = no changes, score 1 = erythema and/or mild swelling, score 2 = severe swelling and/or appearance of deformities, score 3 = immobility due to loss of flexion and extension.

Cloning of the bovine whey gp39 cDNA

A Cow mammary gland Lambda BlueMid(-) cDNA library (oligo dT and random-primed, custom made by Clontech Laboratories Inc.) was amplified using the XL1 Blue MRF strain as a host and library plating and preparation of filter replicas were performed according to the manufacturers instructions. Filters were hybridised with a PvuII fragment from HC gp-39 cDNA after ³²P labelling using an oligolabelling kit (Pharmacia). Positive plaques were amplified, rescreened and insert DNA of the corresponding phages were sequenced using a Thermosequenase kit (Amersham). The sequence is indicated in SEQ ID NO: 3. At nucleotide position 954 one clone contained a C whereas another contained a T.

RESULTS**Arthritogenicity of Bovine 39 kDa whey protein**

Following one injection of 25 µg Bovine 39 kDa whey protein mixed with IFA, all mice gradually developed arthritis (Table 5). Signs of arthritis were observed first at 8-20 days after sensitization in fore- or hind paws. The course of Bovine 39 kDa whey protein-induced arthritis was characterized by relapses occurring periodically in hind paws and gradually developed from

mild arthritis into a more severe arthritis (disease progression was followed for 70 days). Very often a symmetrical distribution of afflicted joints was observed, meaning that both fore paws or both hind paws showed arthritic signs at the same time.

Arthritis induction was similarly achieved using 5, 1 or 0.2 instead of 25 µg Bovine 39 kDa whey protein mixed with IFA.

Mice induced with either 1 or 0.2 µg Bovine 39 kDa whey protein developed a more mild form of arthritis: 7 and 6 mice respectively developed a mild arthritis, whereas only 3 and 4 mice developed severe arthritis.

At the end of the experiment, cumulative scores per scoring day for each group of mice were calculated (Figure 2). Following immunization with 25µg Bovine 39 kDa whey protein, a maximum score of 27 (25 µg) was reached after 30 days, followed by distinct relapses. When lower dosages of Bovine 39 kDa whey protein were used for immunization, animals showed a somewhat lower arthritis score compared with mice induced with the 25µg dose. Although lower scores were achieved, a relapsing pattern of arthritis was found in all groups.

In control, PBS treated mice sometimes a slight and transient swelling of hind paws was observed. This phenomenon was also observed in naive, non-injected animals. This mean cumulative swelling per scoring day for these groups of mice had a mean value of 3.4 and was considered as biological background variation.

In conclusion, Bovine 39 kDa whey protein (both 25, 5, 1 and 0.2 µg) is capable to induce a progressive arthritis in female Balb/c mice. The chronic nature of arthritis induction by Bovine 39 kDa whey protein, characterized by recurrent relapses in addition to symmetrical affliction of joints, is reminiscent of disease progression in rheumatoid arthritis (RA).

Table 5: Initiation and progression of arthritis in Bovine 39 kDa whey protein sensitized Balb/c mice.

sensitization (n=10)	arthritis onset (day)		arthritis signs		
	FP	HP	none	mild	severe
0.2 µg B 39 kDa whey protein	-----, -----,14,18	8,14,14,14,18, 18,18,18,30,54	0	6	4
1 µg B 39 kDa whey protein	-----, 8,12,12,16,18	8,8,8,8,8, 8,14,16,18,18	0	7	3
5 µg B 39 kDa whey protein	-----, 8,10,10,12,14	6,6,8,10,10 16,16,18,20,30	0	0	10
25 µg B 39 kDa whey protein	-----, -----,12	8,8,10,12,12 14,14,18,18,20	0	1	9

FP = fore paw, HP = hind paw. Arthritis signs: none: score = 0, mild: score per animal does not exceed 2, severe: score per animal ≥ 3 .

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: AKZO NOBEL N.V.
(B) STREET: Velperweg 76
(C) CITY: Arnhem
(E) COUNTRY: The Netherlands
(F) POSTAL CODE (ZIP): 6824 BM
(G) TELEPHONE: 0412 666376
(H) TELEFAX: 0412 650592
(I) TELEX: 37503 akpha nl

(ii) TITLE OF INVENTION: Novel peptides derived from autoantigen for use in immunotherapy of autoimmune diseases

(iii) NUMBER OF SEQUENCES: 3

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Tyr Lys Leu Val Cys Tyr Tyr Thr Ser Trp Ser Gln Tyr Arg Glu Gly
1 5 10 15

-23-

Asp Gly Ser Cys Phe Pro Asp Ala Leu Asp Arg Phe Leu Cys Thr His
 20 25 30

Ile Ile Tyr Ser Phe Ala Asn Ile Ser Asn Asp
 35 40

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 383 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Gly Leu Arg Ala Ala Gln Thr Gly Phe Val Val Leu Val Leu Leu
 1 5 10 15

Gln Ser Cys Ala Ala Tyr Lys Leu Ile Cys Tyr Tyr Thr Ser Trp Ser
 20 25 30

Gln Tyr Arg Glu Gly Asp Gly Ser Cys Phe Pro Asp Ala Ile Asp Pro
 35 40 45

Phe Leu Cys Thr His Val Ile Tyr Ser Phe Ala Asn Ile Ser Asn Asn
 50 55 60

Glu Ile Asp Thr Trp Glu Trp Asn Asp Val Thr Leu Tyr Asp Thr Leu
 65 70 75 80

Asn Thr Leu Lys Asn Arg Asn Pro Asn Leu Lys Thr Leu Leu Ser Val
 85 90 95

Gly Gly Trp Asn Phe Gly Ser Gln Arg Phe Ser Lys Ile Ala Ser Lys
 100 105 110

-24-

Thr Arg Ser Arg Arg Thr Phe Ile Lys Ser Val Pro Pro Phe Leu Arg
 115 120 125

Thr His Gly Phe Asp Gly Leu Asp Leu Ala Trp Leu Tyr Pro Gly Trp
 130 135 140

Arg Asp Lys Arg His Leu Thr Thr Leu Val Lys Glu Met Lys Ala Glu
 145 150 155 160

Phe Val Arg Glu Ala Gln Ala Gly Thr Glu Gln Leu Leu Leu Ser Ala
 165 170 175

Ala Val Pro Ala Gly Lys Ile Ala Ile Asp Arg Gly Tyr Asp Ile Ala
 180 185 190

Gln Ile Ser Arg His Leu Asp Phe Ile Ser Leu Leu Thr Tyr Asp Phe
 195 200 205

His Gly Ala Trp Arg Gln Thr Val Gly His His Ser Pro Leu Phe Arg
 210 215 220

Gly Gln Glu Asp Ala Ser Ser Asp Arg Phe Ser Asn Ala Asp Tyr Ala
 225 230 235 240

Val Ser Tyr Met Leu Arg Leu Gly Ala Pro Ala Asn Lys Leu Val Met
 245 250 255

Gly Ile Pro Thr Phe Gly Arg Ser Tyr Thr Leu Ala Ser Ser Lys Thr
 260 265 270

Asp Val Gly Ala Pro Ile Ser Gly Pro Gly Ile Pro Gly Gln Phe Thr
 275 280 285

Lys Glu Lys Gly Ile Leu Ala Tyr Tyr Glu Ile Cys Asp Phe Leu His
 290 295 300

Gly Ala Thr Thr His Arg Phe Arg Asp Gln Gln Val Pro Tyr Ala Thr
 305 310 315 320

Lys Gly Asn Gln Trp Val Ala Tyr Asp Asp Gln Glu Ser Val Lys Asn
 325 330 335

-25-

Lys Ala Arg Tyr Leu Lys Asn Arg Gln Leu Ala Gly Ala Met Val Trp
340 345 350

Ala Leu Asp Leu Asp Asp Phe Arg Gly Thr Phe Cys Gly Gln Asn Leu
355 360 365

Ala Phe Pro Leu Thr Ser Ala Ile Lys Asp Val Leu Ala Glu Val
370 375 380

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1152 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

ATGGGGCTGA GGGCGGCTCA GACAGGTTTT GTGGTCCTGG TGCTGCTCCA GAGCTGTGCT	60
GCATACAAGC TGATCTGCTA CTACACCAGC TGGTCCCAGT ACCGGGAGGG TGATGGGAGC	120
TGCTTCCCAG ACGCCATCGA CCCCTTCCTG TGCACCCATG TCATCTACAG CTTTGCCAAC	180
ATAAGCAACA ATGAGATCGA CACCTGGGAG TGGAATGACG TGACGCTCTA TGACACACTG	240
AACACACTCA AGAACAGGAA CCCCAACCTG AAGACCCTCC TATCTGTTGG AGGATGGAAC	300
TTCGGTTCTC AAAGATTTTC CAAGATAGCT TCCAAGACCC GGAGTCGCAG GACTTTCATC	360
AAGTCGGTGC CACCATTTCT GCGGACCCAT GGCTTTGATG GACTGGACCT AGCATGGCTC	420
TACCCCGGGT GGAGAGACAA GCGGCATCTC ACCACTCTGG TCAAGGAAAT GAAGGCTGAG	480
TTTGTAAGGG AAGCCCAAGC AGGCACAGAG CAGCTTCTGC TCAGTGCAGC AGTACCTGCA	540


GGGAAGATTG CTATTGACAG AGGCTATGAC ATCGCCCAGA TATCCCGACA CCTGGACTTC	600
ATCAGCCTTT TGACCTATGA CTTTCACGGA GCCTGGCGCC AGACAGTCGG ACACCACAGC	660
CCCCTGTTTC GAGGCCAGGA AGATGCAAGT TCTGACAGAT TCAGTAACGC TGA CTACGCT	720
GTGAGCTACA TGCTGAGGCT GGGGGCTCCA GCCAATAAGC TGGTGATGGG TATCCCCACT	780
TTTGGGAGGA GCTACACTCT GGCCTCTTCC AAGACAGATG TGGGAGCCCC CATCTCAGGG	840
CCAGGAATTC CAGGCCAGTT CACCAAGGAG AAAGGGATCC TTGCCTATTA TGAGATCTGT	900
GACTTCCTCC ACGGAGCCAC CACCCACAGA TTCCGTGACC AGCAGGTCCC CTATGCCACC	960
AAGGGCAACC AGTGGGTGGC GTATGACGAC CAGGAGAGTG TCAAAAACAA GGCACGGTAC	1020
CTGAAGAACA GGCAGCTGGC TGGCGCCATG GTGTGGGCCC TGGACTTGGA TGA CTTCGG	1080
GGCACCTTCT GTGGGCAGAA CCTGGCCTTT CCTCTCACAA GTGCCATCAA AGATGTGCTT	1140
GCTGAGGTGT AG	1152

Claims

1. Protein comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence YKLVCYYTSWSQYREGDGSCFPDALDRFLCTHIIYS-
5 FANISND (SEQ ID NO: 1) for use as a therapeutical substance.
2. Use according to claim 1, characterized in that said protein is bovine 39 kDa whey protein or human YKL-39 protein.
3. Pharmaceutical composition comprising one or more proteins comprising an amino acid
10 sequence which exhibits at least 50% homology with the amino acid sequence YKLVCYYTSWSQYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID NO: 1), and a pharmaceutical acceptable carrier.
4. Pharmaceutical composition according to claim 3, characterized in that said protein is bovine 39 kDa whey protein or human YKL-39 protein.
5. Use of the protein comprising an amino acid sequence which exhibits at least 50%
15 homology with the amino acid sequence YKLVCYYTSWSQYREGDGSCFPDALDR-FLCTHIIYSFANISND (SEQ ID NO: 1) for the manufacture of a pharmaceutical preparation for the induction of specific T-cell tolerance to the homologous autoantigen in mammals suffering from T-cell mediated cartilage destruction.
6. Use according to claim 5, characterized in that said protein is bovine 39 kDa whey
20 protein or human YKL-39 protein.
7. One or more proteins comprising an amino acid sequence which exhibits at least 50% homology with the amino acid sequence YKLVCYYTSWSQYREGDGSCFPDALDR-FLCTHIIYSFANISND (SEQ ID NO: 1) for use in a method to induce arthritis in animals, preferably mice.
- 25 8. A method according to claim 7, characterized in that said protein is bovine 39 kDa whey protein or human YKL-39 protein.
9. Animals which suffer from arthritis, whereby said arthritis was induced by the method according to claim 7 or 8.

10. Use of the animals according to claim 9 for use in the screening of novel drugs for the treatment of arthritis, especially rheumatoid arthritis.
11. Protein having the amino acid sequence of SEQ ID NO: 2.
12. Isolated DNA encoding the amino acid sequence according to claim 11.
- 5 13. Isolated DNA according to claim 12 having the DNA sequence of SEQ ID NO: 3.

1/5

FIG. 1A: Arthritis score fore paw
non tolerized

2 / 5




FIG. 1B: Arthritis score fore paw
tolerized


FIG. 1C: Arthritis score hind paw
non tolerized

4 / 5

FIG. 1D: Arthritis score hind paw
tolerized

5/5

FIG. 2: Arthritis score hind paws



SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/EP 97/01903

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/12 C07K14/47 A61K38/17 A01K67/027

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K C12P A61K A01K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section Ch, Week 8810 Derwent Publications Ltd., London, GB; Class B04, AN 88-068419 XP002039030 & JP 63 023 898 A (DAINIPPON PHARM CO LTD) , 1 February 1988 see abstract	1
X	--- DAINIPPON PHARM KK.: "NEW POLYPEPTIDE AND DNA ENCODING IT - RELATED TO PROTECTIVE MECHANISMS SUCH AS IMMUNE RESPONSE " EMBL SEQUENCE DATA LIBRARY, 19 October 1990, HEIDELBERG, GERMANY, XP002039024 ACCESSION No. P81342 --- -/-	1



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

26 September 1997

Date of making of the international search report

16.09.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Holtorf, S

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 97/01903

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X	<p>NYIRKOS, P. AND GOLDS, E.E.: "HUMAN SYNOVIAL CELLS SECRETE A 39kDa PROTEIN SIMILAR TO A BOVINE MAMMARY PROTEIN EXPRESSED DURING THE NON-LACTATING PERIOD" BIOCHEMICAL JOURNAL, vol. 268, 1990, pages 265-268, XP002039020 see figure 3</p>	11-13
X	<p>--- REJMAN, J.J. AND HURLEY, W.L.: "ISOLATION AND CHARACTERIZATION OF A NOVEL 39 KILODALTON WHEY PROTEIN FROM BOVINE MAMMARY SECRETIONS COLLECTED DURING THE NONLACTATING PERIOD" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, vol. 150, no. 1, 1988, pages 329-334, XP002039021 cited in the application see figure 4</p>	11-13
A	<p>--- WO 95 01995 A (UNIV CALIFORNIA ;PRICE PAUL A (US); JOHANSEN JULIA S (DK)) 19 January 1995 cited in the application pages 6,19,30,36 example 1,6,7</p>	1-13
A	<p>--- HAKALA, B.E., ET AL.: "HUMAN CARTILAGE gp-39, A MAJOR SECRETORY PRODUCT OF ARTICULAR CHONDROCYTES AND SYNOVIAL CELLS, IS A MAMMALIAN MEMBER OF A CHITINASE PROTEIN FAMILY" THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 268, no. 34, 5 December 1993, pages 25803-25810, XP002039022 cited in the application see the whole document</p>	1-13
A	<p>--- HAKALA, B.E., ET AL.: "HUMAN CARTILAGE gp-39, A MAJOR SECRETORY PRODUCT OF ARTICULAR CHONDROCYTES AND SYNOVIAL CELLS, IS A MAMMALIAN MEMBER OF A CHITINASE PROTEIN FAMILY" EMBL SEQUENCE DATA LIBRARY, 1 July 1993, HEIDELBERG, GERMANY, XP002039023 ACCESSION No. P36222;P30923</p>	1-13
A	<p>--- WO 95 27500 A (BRIGHAM & WOMENS HOSPITAL) 19 October 1995 page 1-6; examples</p>	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 97/01903

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
A	<p>BOOT, R.G., ET AL . : "CLONING OF A cDNA ENCODING CHITOTRIOSIDASE, A HUMAN CHITINASE PRODUCED BY MACROPHAGES" THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 44, 3 November 1995, pages 26252-26256, XP002039025 cited in the application see page 26255, right-hand column</p> <p style="text-align: center;">---</p>	1-13
A	<p>SHACKELTON, L. M., ET AL . : "IDENTIFICATION OF A 38-kDa HEPARIN-BINDING GLYCOPROTEIN (gp38k) IN DIFFERENTIATING VASCULAR SMOOTH MUSCLE CELLS AS A MEMBER OF A GROUP OF PROTEINS ASSOCIATED WITH TISSUE REMODELING" THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 22, 2 July 1995, pages 13076-13083, XP002039027 cited in the application see the whole document</p> <p style="text-align: center;">---</p>	1-13
A	<p>KINGSLEY, G., ET AL . : "IMMUNOTHERAPY IN RHEUMATIC DISEASE: AN IDEA WHOSE TIME HAS COME - OR GONE ?" IMMUNOLOGY TODAY, vol. 17, no. 1, January 1996, pages 9-12, XP002039028 see the whole document</p> <p style="text-align: center;">---</p>	1-13
L,P, X	<p>WO 96 13517 A (AKZO NOBEL NV ;BOOTS ANNA MARIA HELENA (NL); VERHEIJDEN GIJSBERTUS) 9 May 1996 see the whole document</p> <p style="text-align: center;">---</p>	1,3,5,7, 9,10
T	<p>VERHEIJDEN, G.F.M., ETAL.: "HUMAN CARTILAGE GLYCOPROTEIN-39 AS A CANDIDATE AUTOANTIGEN IN RHEUMATOID ARTHRITIS" ARTHRITIS AND RHEUMATISM, vol. 40, no. 6, June 1997, pages 1115-1125, XP002039029 see the whole document</p> <p style="text-align: center;">-----</p>	1-13

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/01903

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9501995 A	19-01-95	EP 0710251 A JP 9500209 T	08-05-96 07-01-97
WO 9527500 A	19-10-95	AU 2246795 A AU 2277695 A CA 2185353 A CA 2187345 A EP 0752879 A EP 0752880 A HU 74900 A HU 76101 A NO 964199 A NO 964200 A WO 9527499 A	30-10-95 30-10-95 19-10-95 19-10-95 15-01-97 15-01-97 28-02-97 30-06-97 03-10-96 03-10-96 19-10-95
WO 9613517 A	09-05-96	AU 3925295 A EP 0733065 A FI 962619 A HU 74847 A NO 962695 A PL 315198 A	23-05-96 25-09-96 25-06-96 28-02-97 26-06-96 14-10-96



14

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 38/17		A3	(11) International Publication Number: WO 00/04917
			(43) International Publication Date: 3 February 2000 (03.02.00)
(21) International Application Number: PCT/EP99/05331 (22) International Filing Date: 19 July 1999 (19.07.99) (30) Priority Data: 98202471.3 23 July 1998 (23.07.98) EP (71) Applicant (for all designated States except US): AKZO NOBEL N.V. [NL/NL]; Velperweg 76, NL-6824 BM Arnhem (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): MILTENBURG, Andreas, Martinus, Maria [NL/NL]; Stakenborg 19, NL-5346 VH Oss (NL). BOOTS, Anna, Maria, Helena [NL/NL]; Verlengde Torenstraat 10, NL-5366 AV Megen (NL). (74) Agent: HERMANS, F., G., M.; P.O. Box 20, NL-5340 BH Oss (NL).			(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 20 July 2000 (20.07.00)
(54) Title: USE OF HC gp-39 IN IMMUNE DISEASES			
(57) Abstract The present invention relates to the use of HC gp-39 to prevent inflammatory diseases. More specifically, HC gp-39 or fragments thereof can be used to modulate the reactivity of lymphocytes which are reactive to antigens other than HC gp-39 but which are present in the same tissue as where HC gp-39 is being expressed.			



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		



INTERNATIONAL SEARCH REPORT

Int. Application No.

PCT/EP 99/05331

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 805 206 A (SMITHKLINE BEECHAM CORPORATION) 5 November 1997 (1997-11-05) the whole document	1,3
X	VERHEIJDEN G F ET AL: "Human cartilage glycoprotein -39 as a candidate autoantigen in rheumatoid arthritis." ARTHRITIS AND RHEUMATISM, (1997 JUN) 40 (6) 1115-25, XP002089334 cited in the application the whole document	1,3
A	EP 0 823 478 A (SMITHKLINE BEECHAM CORPORATION) 11 February 1998 (1998-02-11) the whole document -/-	1

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

23 March 2000

Date of mailing of the international search report

04/04/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5816 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Moreau, J



INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 99/05331

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 13517 A (AKZO NOBEL) 9 May 1996 (1996-05-09) cited in the application the whole document	1-3
X	WO 97 40149 A (AKZO NOBEL) 30 October 1997 (1997-10-30) page 4 -page 6	1,3
P,X	COENEN DE ROO C J J ET AL: "Human cartilage GP-39 induces and modifies a chronic relapsing arthritis in mice." 62ND NATIONAL SCIENTIFIC MEETING OF THE AMERICAN COLLEGE OF RHEUMATOLOGY AND THE 33RD NATIONAL SCIENTIFIC MEETING OF THE ASSOCIATION OF RHEUMATOLOGY HEALTH PROFESSIONALS;SAN DIEGO, CALIFORNIA, USA; NOVEMBER 8-12, 1998, vol. 41, no. 9 SUPPL., 1998, page S214 XP002133815 Arthritis & Rheumatism Sept., 1998 ISSN: 0004-3591 the whole document	1-3



INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Patent Application No

PCT/EP 99/05331

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 805206	A	05-11-1997	JP 10066585 A	10-03-1998
EP 823478	A	11-02-1998	US 5811535 A	22-09-1998
			JP 10179170 A	07-07-1998
WO 9613517	A	09-05-1996	AT 177756 T	15-04-1999
			AU 696827 B	17-09-1998
			AU 3925295 A	23-05-1996
			BR 9506377 A	16-09-1997
			CN 1168677 A	24-12-1997
			DE 69508380 D	22-04-1999
			DE 69508380 T	09-09-1999
			EP 0733065 A	25-09-1996
			ES 2130672 T	01-07-1999
			FI 962619 A	25-06-1996
			GR 3030296 T	30-09-1999
			HU 74847 A	28-02-1997
			JP 9507861 T	12-08-1997
			NO 962695 A	26-06-1996
			NZ 295659 A	26-02-1998
			PL 315198 A	14-10-1996
			TR 960361 A	21-06-1996
			US 5736507 A	07-04-1998
WO 9740149	A	30-10-1997	US 5843449 A	01-12-1998
			AU 2386997 A	12-11-1997
			BR 9708714 A	03-08-1999
			CA 2251584 A	30-10-1997
			CN 1216582 A	12-05-1999
			CZ 9803333 A	17-03-1999
			EP 0904369 A	31-03-1999
			NO 984835 A	16-12-1998
			PL 329350 A	29-03-1999

